

# Zebrafish interactions offer help in studying social behavior disorders

26 July 2018



Two zebrafish engage in a get-to-know-you swimming pattern after being introduced to each other for the first time. A breakdown in social behavior by either fish will alter how they relate to one another, resembling similar disruptions seen by people with some socialization disorders. Credit: University of Oregon

July 26, 2018—University of Oregon scientists have identified brain cells vital to how zebrafish socialize. When the neurons are disabled, their orientation to one another breaks down in ways similar to socialization problems seen in humans with autism spectrum disorders and schizophrenia.

The neurons, which have equivalent genes in mice and in humans, provide a new avenue for research on the basic biological underpinnings of [social behavior](#), said co-author Philip Washbourne, a professor of biology and member of the UO Institute of Neuroscience who has studied autism genes for almost two decades.

The discovery is detailed in a paper published online July 26 by the journal *Current Biology*.

Early in the National Institutes of Health-funded study, lead author Sarah J. Stednitz, a doctoral student, noticed a regular pattern of behavior

between two fish interacting socially. When placed in adjacent tanks they engaged in a get-to-know-you swimming pattern, despite the presence of a clear dividing panel between the fish.

The observation prompted the six-member research team to apply a drug known to impair the function of neurons needed for social interactions in mice. Expecting to see behavioral disruptions in the drugged fish, which the researchers did, they also saw reduced orienting behaviors by the healthy fish.

As the project progressed, fish were exposed to non-fish objects, which failed to initiate social orienting. Eventually, the research team concluded that zebrafish copy each other's motions and are dependent on another socially engaged fish to begin social orienting. The presence of another fish that is not socially engaged is insufficient to trigger interaction regardless of their proximity, a finding that mirrors reciprocal social interaction in humans.

The neurons involved in the study are in a region of the fish brain that is theorized to be the evolutionary and functional equivalent for the regulation of memory, emotion and social behavior in mammals. The research team noted that the findings "are promising for future studies into the visual circuitry required to drive social behavior."

The study came amid a shift in the approach to autism-related research in Washbourne's lab.

"We've been studying genes associated with autism and how they might work at the synaptic level in zebrafish," Washbourne said. "We'd done a lot of that in the spinal cord, but it wasn't very satisfying, because, in the end, people with autism don't have deficits in very simple reflexes.

"I wanted to find a way to get at more complex behavior that might be affected by mutations in autism genes," he said. "We thought of social

behavior, which probably uses the kind of synapses that are affected in autism."

Initially, Stednitz set out to explore basic social behavior in zebrafish using an approach used by other scientists to study shoaling in fish.

"Zebrafish are social animals," Washbourne said. "They like to hang out together in groups. They shoal, they don't school unless threatened."

Shoaling refers to how fish interact in proximity with others, as opposed to schooling where fish swim in groups moving in the same direction.

For the research, the UO team developed novel software to help measure the orientation of fish during social behaviors of some 500 zebrafish.

Definitively making a jump for human implications of the research is a long way off, but the prospect is intriguing based on the study," Washbourne said. "For now, what we're trying to do is make a model of [autism](#)," he said. "What we did in this study doesn't make a model."

In her subsequent work, Stednitz is working with mutated genes in zebrafish that may be the equivalent to mutated genes in humans to see if the fish have deficits in social behavior and are, in essence, autistic.

"If and when we have a mutant [zebrafish](#) that has deficits in social behavior, then we can ask, what can we give that [fish](#) to improve its social interaction? That could lead to drug screens and testing," Washbourne said.

**More information:** Sarah J. Stednitz et al, Forebrain Control of Behaviorally Driven Social Orienting in Zebrafish, *Current Biology* (2018). [DOI: 10.1016/j.cub.2018.06.016](https://doi.org/10.1016/j.cub.2018.06.016)

Provided by University of Oregon

APA citation: Zebrafish interactions offer help in studying social behavior disorders (2018, July 26) retrieved 22 October 2019 from <https://phys.org/news/2018-07-zebrafish-interactions-social-behavior-disorders.html>

*This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.*