


New possibilities for using ozonized erythrocyte mass explored

July 27 2018



Pixiz

The structure of the left ventricular myocardium of animals on the 5th day post-transfusion period in the control series. Credit: Lobachevsky University


Any bleeding results in a decrease of the amount of circulating blood, and the disruption of the adequate supply of tissues with oxygen can lead to death. An important measure aimed at correcting the pathological effects of acute blood loss is to restore the globular volume of blood. However, transfusion with erythrocyte mass is not always effective, since after prolonged storage, it can cause additional sludging and thrombosis of the microcirculatory bed, and as a result, deterioration of the blood gas transport function.

It follows from the research of morphological and functional characteristics of preserved [red blood cells](#) that they need to be rehabilitated, and that there is some potential for such morphological and functional rehabilitation. Lobachevsky University (UNN) researchers focused their attention on ozone therapy as a medical technology that can be successfully applied in the field of physiotherapy of pathological conditions accompanied by tissue hypoxia.

Initially, ozone was applied in medical practice as an antiseptic. It was used to treat poorly healing wounds, pressure sores, gangrene, severe burns and to stop severe bleeding. However, the interest in this method, which proved effective in practice, diminished with the advent of potent new antibiotics. In the 1970s, when it became obvious that the use of antibiotics in some cases was inexpedient, the medicinal properties of ozone reemerged on the international research agenda.

Ozone's anti-hypoxic and antioxidant properties, including those that affect morphological and functional properties of erythrocytes, were demonstrated in numerous studies. This led the scientists at the

Lobachevsky University to assume ozone's high potential in correcting the oxygen transport function of the erythrocyte [mass](#) that deteriorated during storage.



Pixiz

The structure of the left ventricular myocardium of animals of the experimental series on the 5th day post-transfusion period. Credit: Lobachevsky University

"It is important to study not only the state of erythrocytes that are treated with ozone, but also to examine the changes in the state of internal organs when using the ozonized erythrocyte mass. Since the decrease in the volume of circulating blood is associated with a change in hemodynamics, which directly involves the heart, maintaining its stable pumping and contractile function becomes one of the main tasks in case of blood loss," notes Anna Deryugina, head of the Physiology and Anatomy Department at the UNN Institute of Biology and Biomedicine.

UNN researchers conducted a study of the morphological and functional state of erythrocytes and myocardial state in rats with blood loss during the posttransfusion period using ozonated erythrocyte mass. It was shown that the use of ozonated erythrocyte mass increases the electronegativity of erythrocyte membranes, reduces their aggregation, and increases the intensity of metabolic processes in erythrocytes. By compensating acute blood loss with ozonated erythrocyte mass, it is possible to reduce the degree of manifestations of tissue hypoxia, as the rheological properties of [blood](#) and microcirculation of the myocardium improve during the posttransfusion period.

"During the posttransfusion period, myocardial microcirculation enhanced through the use of ozone limits the damage to the architectonics of the microcirculation bed and cardiomyocytes in the myocardium and facilitates the rapid and total recovery of the structural integrity of heart cell organelles. Our research shows that [ozone](#) treatment of erythrocyte mass for transfusion has a cardioprotective effect," concludes Anna Deryugina.

More information: A. V. Deryugina et al, Correction of Metabolic Indicators of Erythrocytes and Myocardium Structure with Ozonized Red Blood-Cell Mass, *Cell and Tissue Biology* (2018). [DOI: 10.1134/s1990519x18030033](https://doi.org/10.1134/s1990519x18030033)

Provided by Lobachevsky University

Citation: New possibilities for using ozonized erythrocyte mass explored (2018, July 27)
retrieved 22 September 2024 from
<https://phys.org/news/2018-07-possibilities-ozonized-erythrocyte-mass-explored.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.