


# The origin of off-taste in onions

15 August 2018



Credit: American Chemical Society

Chopping onions is usually associated with watery and stinging eyes. But after the onions are diced and the tears are dried, the vegetable pieces can sometimes develop an unpleasant bitter taste. Now, one group reports in ACS' *Journal of Agricultural and Food Chemistry* that they have identified previously unknown compounds causing this off-taste.

Thanks to onions' characteristic [taste](#) and strong smell, they are one of the most popular vegetables around the world. Onions are known for their versatility—people frequently eat them raw or add them to numerous dishes as they cook. But as onions are chopped and processed for cooking, a bitter taste that could affect the taste of the resulting meal can sometimes develop. Very little is known about why this particular phenomenon occurs. Previous research has shown that onions release a compound called lachrymatory factor upon being cut that causes eyes to water and sting. In addition, scientists have also concluded that several sulfur-based [compounds](#) are also formed when the [onion](#) is chopped, but none of these caused a bitter taste. So Roman Kubec and colleagues wanted to identify the bitter onion compounds.

The team processed onions with a kitchen juicer. Freshly obtained juice was not bitter, but after 30 minutes, it developed a strong [bitter taste](#). The group performed sensory-guided high-performance liquid chromatography to identify the compounds that formed over time in the onion juice. The

researchers discovered nine groups of new sulfur compounds in the onion [juice](#), which they dubbed allithiolanes. These compounds form spontaneously when the onion is damaged. The allithiolanes were not limited to onions; further testing showed they were also present in leeks, and one of the groups was found in garlic.

**More information:** Roman Kubec et al.

Allithiolanes: Nine Groups of a Newly Discovered Family of Sulfur Compounds Responsible for the Bitter Off-Taste of Processed Onion, *Journal of Agricultural and Food Chemistry* (2018). DOI: [10.1021/acs.jafc.8b03118](https://doi.org/10.1021/acs.jafc.8b03118)

## Abstract

The compounds responsible for the bitter off-taste of processed onion (*Allium cepa*) were studied. Using a series of sensory-guided HPLC fractionations, the existence of nine groups of hitherto unknown sulfur compounds has been revealed. On the basis of spectroscopic data (MS, NMR, and IR), it was found that these compounds, trivially named allithiolanes A–I, are members of a large family of structurally closely related derivatives of 3,4-dimethylthiolane S-oxide, with the general formulas of  $C_xH_yO_2S_4$ ,  $C_xH_yO_3S_5$ , and  $C_xH_yO_4S_6$  ( $x = 10–18$ ,  $y = 18–30$ ). The presence of multiple stereoisomers was observed for each group of allithiolanes. Allithiolanes possess an unpleasantly bitter taste with detection thresholds in the range of 15–30 ppm. Formation pathways of these newly discovered sulfur compounds were proposed.

Provided by American Chemical Society

APA citation: The origin of off-taste in onions (2018, August 15) retrieved 5 December 2020 from <https://phys.org/news/2018-08-off-taste-onions.html>

*This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.*