

Fiber optic sensor measures tiny magnetic fields

19 September 2018

Researchers have developed a light-based technique for measuring very weak magnetic fields, such as those produced when neurons fire in the brain. The inexpensive and compact sensors could offer an alternative to the magnetic resonance imaging (MRI) systems currently used to map brain activity without the expensive cooling or electromagnetic shielding required by MRI machines.

"A portable, low-cost brain imaging system that can operate at room temperature in unshielded environments would allow real-time brain activity mapping after potential concussions on the sports [field](#) and in conflict zones where the effect of explosives on the brain can be catastrophic," said researcher member Babak Amirsolaimani of the University of Arizona, Tucson.

As detailed in The Optical Society (OSA) journal *Optics Letters*, the researchers fabricated the magnetic [sensors](#) using optical fibers and a newly developed polymer-nanoparticle composite that is sensitive to magnetic fields. The sensors can detect the brain's [magnetic field](#), which is 100 million times weaker than the magnetic field of earth.

The researchers also showed that the new sensor can detect the weak magnetic pattern of a human heartbeat and has the capability to detect magnetic fluctuations that change every microsecond from an area as small as 100 square microns.

"The all-optical design of the sensor means it could be fabricated inexpensively on a silicon photonics chip, making it possible to produce a system that is almost as small as the sensor's 10-micron-diameter optical fiber," said Amirsolaimani. "Multiple sensors could then be used together to provide high spatial resolution brain mapping."

The new sensors could help scientists better understand the activity of the brain and diseases of

the brain such as dementia and Alzheimer's. They might also be useful for measuring the magnetic fields used to predict volcanic eruptions and earthquakes, identify oil and minerals for excavation and detect military submarines.

Optical detection of magnetic fields

The optical method for detecting weak magnetic fields takes advantage of the fact that a magnetic field causes the polarization of light to rotate, with the degree of rotation dependent on the material through which the light passes. The researchers developed a new composite material made of nanoparticles dispersed in a polymer that imparts a detectable polarization rotation in light when very weak magnetic fields are present.

They selected nanoparticles based on magnetite and cobalt because these materials exhibit very high magnetic sensitivity. They then optimized the size, spacing and coating of the nanoparticles to create a composite material that is extremely sensitive to magnetic fields.

The researchers detected the polarization rotation using an optical interferometer. This works by splitting laser light into two paths, one of which passes through the highly-sensitive material while the other does not. The polarization of each light path is detected and compared to measure fluctuations in very small magnetic fields.

When detecting weak magnetic fields, noise can easily cover up the signal being detected. For this reason, the researchers used an interferometer setup that eliminates ambient environmental effects such as vibration and temperature fluctuations. This setup kept noise levels very close to the theoretical limit of the optical design, which was key for detecting very [weak magnetic fields](#).

The researchers used the sensors to measure the magnetic field created by electrical impulses

produced during the human heartbeat. They were able to detect a clear magnetic signal exhibiting high contrast, demonstrating the technology's potential as a simple replacement for electrocardiography, or ECG, tests commonly performed to detect heart problems.

Next, the researchers plan to study the long-term stability of the sensors and how well they withstand environmental changes. They also want to fabricate several hundred sensors to make a system for evaluating and imaging the entire magnetic field of a human brain.

More information: Babak Amirsolaimani et al, High sensitivity magnetometer using nanocomposite polymers with large magneto-optic response, *Optics Letters* (2018). DOI: [10.1364/OL.43.004615](https://doi.org/10.1364/OL.43.004615)

Provided by Optical Society of America

APA citation: Fiber optic sensor measures tiny magnetic fields (2018, September 19) retrieved 19 October 2019 from <https://phys.org/news/2018-09-fiber-optic-sensor-tiny-magnetic.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.