

# Four newly discovered Milky Way neighbors

1 October 2018



A Magellan megacam optical image of the dwarf galaxy Phoenix II. Yes, the faint galaxy's stars are indeed hard to spot amongst the many foreground stars of the Milky Way. A new paper provides sensitive new observations and constraints on four of the Milky Way's dwarf galaxy neighbors. Credit: Mutlu-Pakdil et al. 2018

Ultra-faint dwarf galaxies are the smallest, most dark matter dominated, and least chemically enriched stellar systems in the universe, and are important targets for understanding dark matter and galaxy formation. They comprise by number the majority of galaxies in the universe, and not least, dwarf galaxies around the Milky Way provide crucial empirical input for verifying formation scenarios of our own galaxy. There are currently about sixty dwarf galaxies associated with the Milky Way and closer than about one million light-years; the Andromeda Galaxy, our closest large neighbor spiral galaxy, is two and one-half million light-years away.

Many new MW satellite [galaxies](#) have been

discovered in the last few years, but some have been called into question by more sensitive imaging campaigns and most have only poorly constrained properties. CfA astronomer Nelson Caldwell was a member of a team that used the Magellan Clay telescope and the Megacam instrument to obtain images of four nearby [dwarf galaxies](#) probing nearly sixteen times fainter than previous measurements. The images reveal new stars and other objects, including extended structures, and enabled the astronomers to revise key parameters of these galaxies.

One of the dwarfs, Sagittarius II, with a gas mass of only 1300 solar-masses, is unusual in that it is small in size even for a dwarf galaxy and might instead be considered as the most extended globular cluster of stars for its brightness. Another, Reticulum II, is the most elongated dwarf galaxy known (nearly eight times longer than it is wide). A third, Tucana III, seems to be associated with a stream of material flowing into the MW and may be tidally disrupted. The sensitive new results were unable to measure any gas in any of the objects, but they set new limits, and will help astronomers make a more complete census of the Milky Way's family of galaxies.

**More information:** Burçin Mutlu-Pakdil et al. A Deeper Look at the New Milky Way Satellites: Sagittarius II, Reticulum II, Phoenix II, and Tucana III, *The Astrophysical Journal* (2018). [DOI: 10.3847/1538-4357/aacd0e](https://doi.org/10.3847/1538-4357/aacd0e)

Provided by Harvard-Smithsonian Center for Astrophysics

APA citation: Four newly discovered Milky Way neighbors (2018, October 1) retrieved 22 September 2020 from <https://phys.org/news/2018-10-newly-milky-neighbors.html>

*This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.*