

New research identifies two types of drought across China and how they evolve

17 October 2018



Credit: CC0 Public Domain

Flash drought is a rapidly intensifying water deficit process accompanied by high temperatures in a short period of time. Recently, heat extremes have become more frequent in a warming climate, and have substantially increased the occurrence of flash drought, which threatens crop yields and water supply.

Dr. Linying Wang and Professor Xing Yuan, from the Institute of Atmospheric Physics, Chinese Academy of Sciences, used in-situ observations and reanalysis datasets to explore the long-term variability and trends of two types of flash drought. Their findings are published in *Advances in Atmospheric Sciences*.

Type I flash drought is high-temperature driven, combined with increased evapotranspiration (ET) and decreased soil moisture, while type II flash drought is initialized by a lack of rainfall and combined with decreased ET and warmer temperatures.

They found that both types of flash drought have increased significantly in the past 30 years, but

with a two-fold greater increase for Type I compared with Type II.

They further examined the spatial distribution characteristics of flash drought under two different physical mechanisms and the connection with seasonal drought, and uncovered some interesting findings.

As explained by Dr. Linying Wang, "The anticyclonic circulation patterns can provide favorable conditions for both types of flash drought. But, as the local climate, vegetation and [soil moisture conditions](#) are quite different, the spatial distributions of the two types of flash drought are different as well. In humid and semi-humid regions, where the moisture supply is sufficient, ET is mainly restricted by available energy, so high temperatures can elevate the ET quickly, which explains why Type I flash drought occurs preferentially over humid and semi-humid regions, such as southern China."

"Composite analysis showed that anomalously [high temperatures](#) one pentad before the onset of flash drought leads to a rapid increase in ET and reduced soil moisture. For flash drought associated with seasonal drought, there is a greater likelihood of occurrence during the onset phase of seasonal drought over southern China. In northern China, meanwhile, due to the limited terrestrial moisture supply, ET decreases with the decrease in [soil moisture](#), resulting in Type II flash drought. Also, they are more common in both the onset and recovery phases of seasonal drought", she added.

Professor Xing Yuan further explains that the perfect conditions for flash drought are during dry/wet transition periods, which may provide a basis for the early warning of flash drought by connecting multiscale [drought](#) phenomena.

More information: Linying Wang et al, Two Types of Flash Drought and Their Connections with

Seasonal Drought, *Advances in Atmospheric Sciences* (2018). DOI: [10.1007/s00376-018-8047-0](https://doi.org/10.1007/s00376-018-8047-0)

Provided by Chinese Academy of Sciences

APA citation: New research identifies two types of drought across China and how they evolve (2018, October 17) retrieved 18 May 2021 from <https://phys.org/news/2018-10-drought-china-evolve.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.