


Magnetic fields may be the key to black hole activity

17 October 2018




Artist's conception of the core of Cygnus A, including the dusty donut-shaped surroundings, called a torus, and jets launching from its center. Magnetic fields are illustrated trapping the dust in the torus. These magnetic fields could be helping power the black hole hidden in the galaxy's core by confining the dust in the torus and keeping it close enough to be gobbled up by the hungry black hole. Credit: NASA/SOFIA/Lynette Cook

The [unified model](#), which attempts to explain the different properties of active galaxies, states that the core is surrounded by a donut-shaped dust cloud, called a torus. How this obscuring structure is created and sustained has never been clear, but these new results from SOFIA indicate that magnetic fields may be responsible for keeping the dust close enough to be devoured by the hungry black hole. In fact, one of the fundamental differences between active galaxies like Cygnus A and their less active cousins, like our own Milky Way, may be the presence or absence of a [strong magnetic field](#) around the black hole.

Although celestial magnetic fields are notoriously difficult to observe, astronomers have used polarized light—optical light from scattering and radio light from accelerating electrons—to study magnetic fields in galaxies. But optical wavelengths are too short and the radio wavelengths are too long to observe the torus directly. The infrared wavelengths observed by SOFIA are just right, allowing scientists, for the first time, to target and isolate the dusty torus.

Collimated jets provide astronomers with some of the most powerful evidence that a supermassive black hole lurks in the heart of most galaxies. Some of these black holes appear to be active, gobbling up material from their surroundings and launching jets at ultra-high speeds, while others are quiescent, even dormant. Why are some black holes feasting and others starving? Recent observations from the Stratospheric Observatory for Infrared Astronomy, or SOFIA, are shedding light on this question.

SOFIA data indicate that magnetic fields are trapping and confining dust near the center of the active galaxy, Cygnus A, and feeding material onto the supermassive black hole at its center.



Two images of Cygnus A layered over each other to

show the galaxy's jets glowing with radio radiation (shown in red). Quiescent galaxies, like our own Milky Way, do not have jets like this, which may be related to magnetic fields. The yellow image shows background stars and the center of the galaxy shrouded in dust when observed with visible light. The area SOFIA observed is inside the small red dot in the center. Credit: Optical Image: NASA/STSiC Radio Image: NSF/NRAO/AUI/VLA

SOFIA's new instrument, the High-resolution Airborne Wideband Camera-plus (HAWC+), is especially sensitive to the infrared emission from aligned dust grains. This has proven to be a powerful technique to study magnetic fields and test a fundamental prediction of the unified model: the role of the dusty torus in the active-galaxy phenomena.

"It's always exciting to discover something completely new," noted Enrique Lopez-Rodriguez, a scientist at the SOFIA Science Center, and the lead author on the report of this new discovery. "These observations from HAWC+ are unique. They show us how infrared polarization can contribute to the study of galaxies."

Recent observations of the heart of Cygnus A made with HAWC+ show infrared radiation dominated by a well-aligned dusty structure. Combining these results with archival data from the Herschel Space Observatory, the Hubble Space Telescope and the Gran Telescopio Canarias, the research team found that this powerful active galaxy, with its iconic large-scale jets, is able to confine the obscuring torus that feeds the supermassive black hole using a strong [magnetic field](#).

The results of this study were published in the July 10th issue of *The Astrophysical Journal Letters*.

Cygnus A is in the perfect location to learn about the role magnetic fields play in confining the dusty torus and channeling material onto the supermassive black hole because it is the closest and most powerful active galaxy. More observations of different types of galaxies are necessary to get the full picture of how magnetic fields affect the evolution of the environment

surrounding [supermassive black holes](#). If, for example, HAWC+ reveals highly polarized [infrared emission](#) from the centers of active galaxies but not from quiescent [galaxies](#), it would support the idea that magnetic fields regulate black hole feeding and reinforce astronomers' confidence in the unified model of [active galaxies](#).

More information: Enrique Lopez-Rodriguez et al. The Highly Polarized Dusty Emission Core of Cygnus A, *The Astrophysical Journal* (2018). DOI: [10.3847/2041-8213/aacff5](https://doi.org/10.3847/2041-8213/aacff5)

Provided by NASA

APA citation: Magnetic fields may be the key to black hole activity (2018, October 17) retrieved 1 October 2020 from <https://phys.org/news/2018-10-magnetic-fields-key-black-hole.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.