

Symbiosis a driver of truffle diversity

November 14 2018



The Périgord black truffle (*Tuber melanosporum*) was the first truffle to have its genome sequenced. Its genome was used in a comparative analysis of truffle-forming fungal species reported in *Nature Ecology & Evolution* from an international team including JGI researchers. Credit: Francis Martin

While the sight of black or white truffle being shaved over on pasta is generally considered a sign of dining extravagance, they play an

important role in soil ecosystem services. Truffles are the fruiting bodies of the ectomycorrhizal (ECM) fungal symbionts residing on host plant roots. In many *Ascomycota* and *Basidiomycota* lineages, truffle-forming species have evolved independently in nearly every major group. This suggests that symbiosis drives evolution of truffle diversity and selects for specific traits.

A team led by Francis Martin and his colleagues at the French National Institute for Agricultural Research (INRA), Genoscope, and University of Torino, and including researchers at the U.S. Department of Energy (DOE) Joint Genome Institute (JGI), a DOE Office of Science User Facility, sought insights into the ECM lifestyle of truffle-forming species. As reported in *Nature Ecology & Evolution*, the team conducted a comparative analysis of eight Pezizomycete fungi, including four species prized as delicacies.

A Decade of ECM Symbiosis Studies

The underground networks that link ECM fungi and their host plant roots shape these ecosystems, impacting the plant's health and tolerance to stressors such as drought or disease, as well as the global carbon cycle. Studies have suggested that ectomycorrhizal fungi may slow soil carbon cycling down by limiting nitrogen. The fungi absorb mineral nutrients from the soil and exchange them for sugars that plants produce through photosynthesis.

Through the JGI's Community Science Program, JGI de novo sequenced the genomes of two truffle-forming fungi: the Pig truffle (*Choiromyces venosus*) and, the Desert truffle (*Terfezia boudieri*) as well as of two other Pezizomycetes not forming truffles: *Ascobolus immersus* and *Morchella importuna*. These four genomes as well as the genome of the Piedmont white truffle (*Tuber magnatum*) were annotated through the JGI's pipeline. The genome of the Burgundy Truffle (*T. aestivum*) was

sequenced by Genoscope.

The work builds on earlier fungal symbiosis studies involving the first ECM fungal genome (*Laccaria bicolor*, sequenced a decade ago by the JGI), and the first truffle genome (*T. melanosporum*) sequenced by Genoscope, which is part of this analysis. "We have learned from the *Laccaria bicolor* and *Tuber melanosporum* [genome](#) that the ECM symbiosis evolved by the massive loss of genes involved in plant cell wall degradation (CAZymes) and de novo innovation of communication proteins, such as the mycorrhiza-induced small secreted proteins controlling the host plant immunity," said study senior author and longtime JGI collaborator Martin. In a 2015 study with the JGI, Martin and his colleagues then showed that ECM species evolved from saprotrophs that feed on decaying organic matter, and that these evolutionary patterns are present in many *Basidiomycota* groups.

"Here," Martin added, "we showed that the loss of genes involved in lignocellulose/plant cell wall degradation (CAZymes) and a higher evolution rate of symbiosis-related orphan genes have shaped the genomes of peizomycetes truffles, one of the oldest/basal clade of ectomycorrhizal Ascomycota. This means that similar evolutionary mechanisms have independently driven symbiosis in *Ascomycota* and *Basidiomycota*. We also showed that developmental and metabolic pathways expressed in ectomycorrhizal roots and fruiting bodies of the white (*T. magnatum*) and black (*T. melanosporum*) truffles are unexpectedly very similar, owing to the fact that they diverged ~100 million years ago."

Scenting Next Steps

Detecting and disseminating these underground delicacies depends on trained animals scenting the distinctive odors of truffles. Martin and his team suggest the volatile organic compounds that produce these scents

may also be altered by the microbial communities flourishing into truffles .

'In addition to the small secreted proteins critical for fungal interactions with their plant hosts these small molecules encoded in the [truffle](#) genomes by clusters of secondary metabolite genes offer another channel of communications between fungi and the environment,' adds Igor Grigoriev, the JGI Fungal Program head and concludes: "Thus, the sequenced genomes open doors for exploration of multi-dimensional fungal plant interactions'

More information: Claude Murat et al, Pezizomycetes genomes reveal the molecular basis of ectomycorrhizal truffle lifestyle, *Nature Ecology & Evolution* (2018). [DOI: 10.1038/s41559-018-0710-4](https://doi.org/10.1038/s41559-018-0710-4)

Provided by Lawrence Berkeley National Laboratory

Citation: Symbiosis a driver of truffle diversity (2018, November 14) retrieved 22 May 2024 from <https://phys.org/news/2018-11-symbiosis-driver-truffle-diversity.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.