

Affordable catalyst for carbon dioxide recycling

November 19 2018



Kai Junge Puring, Stefan Piontek and Mathias Smialkowski (from left), members of the group led by Ulf-Peter Apfel, with the electrolysis cell in which the experiments were carried out. Credit: RUB, Marquard

A catalyst for carbon dioxide recycling, mineral pentlandite may also be


a conceivable alternative to expensive precious metal catalysts. This is the result of a study conducted by researchers from Ruhr-Universität Bochum (RUB), Fritz-Haber Institute Berlin and Fraunhofer Umsicht in Oberhausen. Pentlandite had previously been known as a catalyst for hydrogen production. By adding a suitable solvent, the researchers successfully utilised it to convert carbon dioxide into carbon monoxide. The latter is a common source material in the chemical industry.

The research team headed by Dr. Ulf-Peter Apfel, Chair of Inorganic Chemistry I in Bochum, describes the findings together with their colleagues in the journal *Chemical Science* from 5 November 2018.

CO₂ conversion replaces hydrogen production

"The conversion of CO₂ into valuable source materials for the [chemical industry](#) is a promising approach to combatting climate change," says Ulf-Peter Apfel. "However, we currently don't know many cheap and readily available catalysts for CO₂ reduction." Moreover, potentially suitable catalysts primarily facilitate another chemical reaction, i.e. the synthesis of hydrogen—these including pentlandite. Nevertheless, the researchers have successfully converted the mineral to be a CO₂ catalyst.

They generated electrodes from pentlandite and analysed under which conditions production of hydrogen or carbon monoxide took place at their surface. "The decisive factor was [water](#) being present at the electrode surface," summarises Ulf-Peter Apfel. A lot of water shifted the reaction towards [hydrogen production](#), a little water towards carbon monoxide production. By adjusting the water content, the researchers were thus able to generate carbon monoxide and hydrogen mixtures. "Synthetic gas mixtures like this one play a crucial role in the chemical industry," points out Apfel.



The researchers carried out the experiments in this electrolysis cell. Credit: RUB, Marquard

Stable catalyst

Pentlandite consists of iron, nickel and sulphur and resembles catalytically active enzyme centres that occur in nature, such as hydrogen-producing hydrogenases. "A huge advantage of this mineral is the fact that it remains stable when confronted with other chemical compounds that occur in industrial emissions and are poison to many catalysts," explains Apfel.

More information: Stefan Piontek et al, Bio-inspired design: bulk

iron–nickel sulfide allows for efficient solvent-dependent CO₂ reduction, *Chemical Science* (2018). DOI: [10.1039/c8sc03555e](https://doi.org/10.1039/c8sc03555e)

Provided by Ruhr-Universitaet-Bochum

Citation: Affordable catalyst for carbon dioxide recycling (2018, November 19) retrieved 22 September 2024 from <https://phys.org/news/2018-11-catalyst-carbon-dioxide-recycling.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.