


From gamma rays to X-rays: New method pinpoints previously unnoticed pulsar emission

22 November 2018



Observations of PSR J1826-1256 – a radio quiet gamma-ray pulsar – obtained with ESA's XMM-Newton X-ray observatory. Scientists discovered the X-ray pulsed emission from this and two similar sources using a theoretical model that predicts a pulsar's non-thermal X-ray brightness on the basis of its observed gamma-ray brightness. This pulsar has a period of 110.2 milliseconds, appearing alternatively fainter and brighter as its beams of radiation point towards and away from Earth. For illustration purposes, the pulsar's flickering is shown 10 times slower than the actual one in this animated view. Credit: ESA/XMM-Newton/J. Li, DESY, Germany

Based on a new theoretical model, a team of scientists explored the rich data archive of ESA's XMM-Newton and NASA's Chandra space observatories to find pulsating X-ray emission from three sources. The discovery, relying on previous gamma-ray observations of the pulsars, provides a novel tool to investigate the mysterious

mechanisms of pulsar emission, which will be important to understand these fascinating objects and use them for space navigation in the future.

Lighthouses of the Universe, pulsars are fast-rotating neutron stars that emit beams of radiation. As pulsars rotate and the beams alternatively point towards and away from Earth, the source oscillates between brighter and dimmer states, resulting in a signal that appears to 'pulse' every few milliseconds to seconds, with a regularity rivalling even atomic clocks.

Pulsars are the incredibly dense, extremely magnetic, relics of massive stars, and are amongst the most extreme objects in the Universe. Understanding how particles behave in such a [strong magnetic field](#) is fundamental to understanding how matter and magnetic fields interact more generally.

Originally detected through their radio emission, pulsars are now known to also emit other types of radiation, though typically in smaller amounts. Some of this emission is standard [thermal radiation](#) – the type that everything with a temperature above absolute zero emits. Pulsars release thermal radiation when they accrete matter, for example from another star.

But pulsars also emit non-thermal radiation, as is often produced in the most extreme cosmic environments. In pulsars, non-thermal radiation can be created via two processes: synchrotron emission and curvature emission. Both processes involve charged particles being accelerated along magnetic field lines, causing them to radiate light that can vary in wavelength from radio waves to gamma-rays.

Non-thermal X-rays result mostly from synchrotron

emission, while gamma-rays may come from so-called synchro-curvature emission – a combination of the two mechanisms. It is relatively easy to find pulsars that radiate gamma-rays – NASA's Fermi Gamma-Ray Space Telescope has detected more than 200 of them over the past decade, thanks to its ability to scan the whole sky. But only around 20 have been found to pulse in non-thermal X-rays.

"Unlike gamma-ray detecting survey instruments, X-ray telescopes must be told exactly where to point, so we need to provide them with some sort of guidance," says Diego Torres, from the Institute of Space Sciences in Barcelona, Spain.

Aware that there should be many pulsars emitting previously undetected non-thermal X-rays, Torres developed a model that combined synchrotron and curvature radiation to predict whether pulsars detected in gamma-rays could also be expected to appear in X-rays.

"Scientific models describe phenomena that can't be experienced directly," explains Torres.

"This model in particular helps explain the emission processes in pulsars and can be used to predict the X-ray emission that we should observe, based on the known gamma-ray emission."

The model describes the gamma-ray emission of pulsars detected by Fermi – specifically, the brightness observed at different wavelengths – and combines this information with three parameters that determine the pulsar emission. This allows a prediction of their brightness at other wavelengths, for instance in X-rays.

Observed X-ray and gamma-ray emission from three pulsars: J1747-2958 (left), J2021+3651 (centre), and J1826-1256 (right). The X-ray pulsed emission was discovered using a theoretical model that predicts a pulsar's non-thermal X-ray brightness on the basis of its observed gamma-ray brightness. The gamma-ray observations are from NASA's Fermi Gamma-Ray Space Telescope; the X-ray observations are from NASA's Chandra X-ray observatory (left and centre) and ESA's XMM-Newton X-ray observatory (right). The red curve in the graphs represents the best fit of the model describing the overall emission of the sources compared to the observed data (black symbols). In the upper row, the fit was performed using only the gamma-ray data: the value in the X-ray energy range represents the theoretical prediction, which is pretty close to what was later revealed in the observations. In the lower row, the fit includes the X-ray data as well, providing a more accurate description of the phenomenon using the same model. Credit: Adapted from J. Li et al. (2018)


Torres partnered with a team of scientists, led by Jian Li from the Deutsches Elektronen Synchrotron in Zeuthen near Berlin, Germany, to select three known gamma-ray emitting pulsars that they expected, based on the model, to also shine brightly in X-rays. They dug into the data archives of ESA's XMM-Newton and NASA's Chandra X-ray observatories to search for evidence of non-thermal X-ray emission from each of them.

"Not only did we detect X-ray pulsations from all three of the pulsars, but we also found that the spectrum of X-rays was almost the same as predicted by the model," explains Li.

"This means that the model very accurately describes the emission processes within a pulsar."

In particular, XMM-Newton data showed clear X-ray emission from PSR J1826-1256 – a radio quiet gamma-ray pulsar with a period of 110.2 milliseconds. The spectrum of light received from this pulsar was very close to that predicted by the model. X-ray emission from the other two pulsars, which both rotate slightly more quickly, was revealed using Chandra data.

This discovery already represents a significant increase in the total number of pulsars known to



emit non-thermal X-rays. The team expects that many more will be discovered over the next few years as the model can be used to work out where exactly to look for them.

<https://arxiv.org/abs/1811.08339>

Finding more X-ray pulsars is important for revealing their global properties, including population characteristics. A better understanding of pulsars is also essential for potentially taking advantage of their accurate timing signals for future space navigation endeavours.

Provided by European Space Agency

The result is a step towards understanding the relationships between the emission by pulsars in different parts of the electromagnetic spectrum, enabling a robust way to predict the brightness of a pulsar at any given wavelength. This will help us better comprehend the interaction between particles and magnetic fields in pulsars and beyond.

"This [model](#) can make accurate predictions of pulsar X-ray emission, and it can also predict the [emission](#) at other wavelengths, for example visible and ultraviolet," Torres continues.

"In the future, we hope to find new pulsars leading to a better understanding of their global properties."

The study highlights the benefits of XMM-Newton's vast data archive to make new discoveries and showcases the impressive abilities of the mission to detect relatively dim sources. The team is also looking forward to using the next generation of X-ray space telescopes, including ESA's future Athena mission, to find even more pulsars emitting non-thermal X-rays.

"As the flagship of European X-ray astronomy, XMM-Newton is detecting more X-ray sources than any previous satellite. It is amazing to see that it is helping to solve so many cosmic mysteries," concludes Norbert Schartel, XMM-Newton Project Scientist at ESA.

More information: Jian Li et al. Theoretically Motivated Search and Detection of Non-thermal Pulsations from PSRs J1747-2958, J2021+3651, and J1826-1256, *The Astrophysical Journal* (2018). [DOI: 10.3847/2041-8213/aae92b](https://doi.org/10.3847/2041-8213/aae92b) ,

APA citation: From gamma rays to X-rays: New method pinpoints previously unnoticed pulsar emission (2018, November 22) retrieved 24 October 2019 from <https://phys.org/news/2018-11-gamma-rays-x-rays-method-previously.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.