


Classic double-slit experiment in a new light

18 January 2019



contradicts the classical expectation of particle behaviour, but can be explained in quantum-mechanical wave-particle dualism. In contrast, the researchers in Cologne investigated an iridium oxide crystal ($\text{Ba}_3\text{CeIr}_2\text{O}_9$) by means of resonant inelastic X-ray scattering (RIXS).

The crystal is irradiated with strongly collimated, high-energy X-ray photons. The X-rays are scattered by the iridium atoms in the crystal, which take over the role of the slits in Young's classical experiment. Due to the rapid technical development of RIXS and a skilful choice of crystal structure, the physicists were observed the scattering on two adjacent iridium atoms, a so-called dimer.

An intense beam of high-energy X-ray photons (violet) hits two adjacent iridium atoms (green) in the crystal. This excites electrons in the atoms for a short time. The atoms emit X-ray photons which overlap behind the two iridium atoms (red) and can be analyzed as interference images. Credit: Markus Grueninger, University of Cologne

An international research team led by physicists from the University of Cologne has implemented a new variant of the basic double-slit experiment using resonant inelastic X-ray scattering at the European Synchrotron ESRF in Grenoble. This new variant offers a deeper understanding of the electronic structure of solids. Writing in *Science Advances*, the research group have now presented their results in a study titled "Resonant inelastic X-ray incarnation of Young's double-slit experiment."

The double-slit experiment is of fundamental importance in physics. More than 200 years ago, Thomas Young diffracted light at two adjacent slits, thus generating [interference patterns](#) (images based on superposition) behind this double slit. Thus, he demonstrated the wave nature of light. In the 20th century, scientists have shown that electrons or molecules scattered on a double slit show the same [interference pattern](#), which



An international research team has implemented a new variant of the basic double-slit experiment using resonant inelastic X-ray scattering at the European Synchrotron ESRF in Grenoble. Credit: ESRF/Jayet

"The interference pattern tells us a lot about the scattering object, the dimer double slit," says Professor Markus Grueninger, who heads the research group at the University of Cologne. In contrast to the classical [double-slit experiment](#), the inelastically scattered X-ray photons provide information about the excited states of the dimer, in particular their symmetry, and thus about the


dynamic physical properties of the solid.

These RIXS experiments require a modern synchrotron as an extremely brilliant X-ray light source and a sophisticated experimental setup. To specifically excite only the iridium atoms, scientists have to select the very small proportion of photons with the right energy from the broad spectrum of the [synchrotron](#), and the scattered photons are selected even more strictly according to energy and direction of scattering. Only a few photons remain. With the required accuracy, these RIXS experiments are currently only possible at two synchrotrons worldwide, including the ESRF (European Synchrotron Radiation Facility) in Grenoble, where the team from Cologne conducted their experiment.

experiments that will allow us to gain a deeper understanding of the properties and functionalities of solids," says Grueninger.

More information: Resonant inelastic x-ray incarnation of Young's double-slit experiment. *Science Advances* (2019). advances.sciencemag.org/content/5/1/eaav4020

Provided by University of Cologne



The two adjacent iridium atoms (dimer) are shown in green. The elements oxygen (O, red), barium (Ba, grey) and cerium (Ce, turquoise) are also involved in the crystal structure. Credit: Markus Grueninger, University of Cologne

"With our RIXS experiment, we were able to prove a fundamental theoretical prediction from 1994. This opens a new door for a whole series of further

APA citation: Classic double-slit experiment in a new light (2019, January 18) retrieved 26 November 2020 from <https://phys.org/news/2019-01-classic-double-slit.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.