


Scientists discover natural fitness watch in fishes that records their activity levels

January 18 2019



Carbon in the earstone comes from water (blue) & the fish's diet (red). Credit: University of Southampton

An international research team including scientists from the University of Southampton have shown for the first time that the energetic cost of living (the metabolic rate) of fish can be measured in structures that grow in their ears. This new tool can be used to show how fish are influenced by and adapt to changes in their environment, including climate change.

Animals need to eat enough food to provide energy to sustain their bodies, hunt, grow and reproduce. In a changing world, changes to the metabolic cost of life can influence where animals can live. To date, measuring the energetic cost of living in the wild has been difficult, especially for animals that live in [water](#). However, new research has discovered that a [fish](#)'s pace of life is recorded in a small structure in its inner ear, known as an [otolith](#), or earstone.

Earstones help the fish know up from down, determine their speed through the water and aid hearing. The researchers from UK, Norway and Denmark have now shown that it also acts as a fitness watch, revealing whether the fish had an active life or was completely relaxed.

Earstones – the fish fitness watch

The carbon in the earstone originates from two sources: the water the fish lives in and the food it eats to meet its energy demands. The carbon from both sources mix in the blood before some of it ends up in the earstone as [calcium carbonate](#). By carefully analyzing the calcium carbonate, the researchers can determine the how much of the carbon originates from the water or the diet as the two sources have different proportions of the light and the heavy carbon isotopes. When the fish's metabolism increases, for example when it encounters warmer water or is more active, it burns off more food and the proportion of carbon from the food in its earstone increases.

Earstones grow new layers similar to the rings in a tree trunk. Every little increment in the otolith growth reflects the growth of the fish and can be used to document its age. This data can be recovered even after the fish has died. Now, by measuring carbon isotopes in the individual layers, the researchers can tell how the metabolism of a fish has changed on a monthly basis during its life, including how seasonal variations in temperature, food and the fish's behaviour have affected its metabolism.

"Our new results is the key to a treasure chest of new discoveries", says Dr. Clive Trueman, from the University of Southampton, who is one of the co-authors behind the paper that is being published in the journal *Communications Biology* today.

Dr. Ming-Tsung Ching, another co-author from Aarhus University in Denmark added "Now we have the tools to measure and understand how fish are being affected by and adapting to changes in the environment, and how much the fish must eat to gain enough energy to swim, grow and reproduce. This new tool will help us improve our predictions of what happens to the fish including commercially important fish like cod when the environment changes".

Long time series

Otoliths have been studied and stored for over 100 years, so now we can go back in time and see how the metabolism of fish has changed in response to industrial fishing and long-term [climate change](#). But it is not only these scientific collections that contain exciting material for research. It is possible to find very old otoliths in archaeological sites known as middens. These otoliths, millennia old, still contain the [carbon](#) signal that reveals the [metabolic rate](#) of the fish.

"We can use knowledge about ancient fish metabolism to learn about the ability of the fish to adapt to environmental conditions that may come to exist in the near future, or how fish respond to changes in water temperature or food availability. What happens to the fish when the waters become warmer than they are now?", says Prof. Grønkjaer from Aarhus University in Denmark.

Dr. Trueman added: "We have already started looking into the metabolic rates of cod living in the cold, but warming waters, around Greenland. We have otoliths from fish in that region dating back to 1926 which will

help us understand the impact of climate on the well-being of these fish.

"Otoliths are like the black box of fish, they contain an enormous library of information written in a language that we are now starting to decipher."

More information: Ming-Tsung Chung et al, Field metabolic rates of teleost fishes are recorded in otolith carbonate, *Communications Biology* (2019). [dx.doi.org/10.1038/s42003-018-0266-5](https://doi.org/10.1038/s42003-018-0266-5)

Provided by University of Southampton

Citation: Scientists discover natural fitness watch in fishes that records their activity levels (2019, January 18) retrieved 26 April 2024 from <https://phys.org/news/2019-01-scientists-natural-fishes.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.