

Research reveals new species are evolving fastest in Antarctica

January 24 2019, by Anastasia Casagrande



Dr. Tim O'Hara in Marine Vertebrate research laboratory holding a specimen jar containing brittle stars. Credit: Museums Victoria

New research published in *Nature* overturns previous theories about how the stunning biodiversity of the oceans evolved, with important


implications for conservation.

The study, titled "Contrasting processes drive ophiuroid phylodiversity across shallow and deep seafloors," was led by first author Dr. Tim O'Hara, senior curator of marine invertebrates at Museums Victoria.

Biologists have long speculated that evolution is sped up by relatively high tropical temperatures, with development being slower in cooler and [deeper waters](#). However, this research finds that evolution does not follow one course, but rather depends on the geological, climatic and biological history of each ecosystem. Evolution proceeded differently in shallow and deep seas.

Speciation was found to be highest in the coldest region: Antarctica. These waters are still apparently recovering from extinction events of tens of millions of years ago, when ice sheets began to dominate and [water temperatures](#) plummeted. New species that evolved as a result are still in the process of diversifying, and are doing so rapidly.

By contrast, although diversity in tropical deep seas (deeper than 200 metres) is high, it is not an environment that is rapidly producing new species, but rather accumulated its rich biodiversity over millions of years. Tropical deep seas are a refuge for ancient fauna, or "living fossils," mainly due to relatively stable conditions over time.




Ophiuroidea, brittle star. Collected and photographed during the Museums Victoria and partners 'Sampling the Abyss' voyage on the RV Investigator. Credit: Museums Victoria

To study patterns of evolution across the world's oceans, the team focused on the evolution of deep-sea [brittle stars](#) (Ophiuroidea). These strange, spiny echinoderms with a typically circular body and five long, flexible arms, are abundant on the seafloor globally. Although they will be unfamiliar to many, their abundance makes them the perfect group for studying large-scale patterns of how marine life arose and spread around the planet.

The researchers utilised data collected on 2017's pioneering "Sampling the Abyss" voyage aboard CSIRO Marine National Facility research

vessel Investigator, led by Museums Victoria. The month-long expedition explored the abyssal ocean depths off the eastern coast of Australia for the first time. Dr. O'Hara was Chief Scientist on the voyage, and this publication is the first major paper to be published as a result of the voyage.

DNA was used to reconstruct a comprehensive picture of how brittle stars have evolved across the Indian and Pacific Oceans in the southern hemisphere. Dr. O'Hara explained, "Museum collections are a treasure house of preserved biodiversity collected from thousands of scientific expeditions. Sequencing the DNA from these specimens can unlock the history of life on our planet. The digitisation and DNA sequencing of [museum collections](#) is providing a new way of looking at how life has evolved and spread around the globe."



Map showing geographic distribution of samples. Credit: Museums Victoria

The deep sea is the world's largest ecosystem, an ancient ark of relics from the dinosaur era, where "living fossils" exist at the same time that new species are fast evolving. These environments require as much protection as more famous and familiar habitats, like coral reefs and mangroves. Yet a lack of knowledge about [marine life](#) in these dark waters has made it unclear how best to protect and preserve these environments from human exploitation like fishing or deep-sea mining.

Dr. O'Hara and his team's paper is the result of what he hopes will be the first stage of a global project to shed further light on processes of [evolution](#) in precious deep sea environments, and how we can best protect them.

More information: Timothy D. O'Hara et al. Contrasting processes drive ophiuroid phylodiversity across shallow and deep seafloors, *Nature* (2019). [DOI: 10.1038/s41586-019-0886-z](https://doi.org/10.1038/s41586-019-0886-z)

Provided by Museums Victoria

Citation: Research reveals new species are evolving fastest in Antarctica (2019, January 24) retrieved 19 September 2024 from <https://phys.org/news/2019-01-reveals-species-evolving-fastest-antarctica.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.