

For some whales, sonar may provoke suicidal behaviour: study

January 30 2019, by Marlowe Hood



Beaked whales get stressed by sonar and can suffer decompression like scuba divers, according to researchers

Scientists have long known that some beaked whales beach themselves and die in agony after exposure to naval sonar, and now they know why: the giant sea mammals suffer decompression sickness, just like scuba

divers.

At first blush, the explanation laid out Wednesday by 21 experts in the *Proceedings of the Royal Society B* seems implausible.

Millions of years of evolution have turned whales into perfectly calibrated diving machines that plunge kilometres (miles) below the surface for hours at a stretch, foraging for food in the inky depths.

The heart rate slows, blood flow is restricted, oxygen is conserved.

So how could the ocean's most accomplished deep-sea diver wind up with nitrogen bubbles poisoning its veins, like a scuba novice rising too quickly to the surface?

Short answer: beaked whales—especially one species known as Cuvier's—get really, really scared.

"In the presence of sonar they are stressed and swim vigorously away from the sound source, changing their diving pattern," lead author Yara Bernaldo de Quiros, a researcher at the Institute of Animal Health at the University of Las Palmas de Gran Canaria, Spain, told AFP.

"The [stress response](#), in other words, overrides the diving response, which makes the animals accumulate nitrogen," she added. "It's like an adrenalin shot."

One type of sonar in particular throws these whales off balance.

Cuvier's beaked whale

Most commonly spotted of the beaked whales, a group of species known as deep-divers that capture their prey by sucking them into their mouths



Max length:
7 m

Ziphius cavirostris
Status: least concerned

**Global population estimate:
up to 100,000**

Found in almost all sub-polar, temperate, sub-tropical and tropical waters, generally where the depth is more than 1,000 m

*Mediterranean
subpopulation*

Status: vulnerable
Population: 5,000 - 7,000

**Maximum dive depth and duration
reliably recorded at over 1,800m
and up to 85 mins, using
echolocation to hunt for squid, fish
and crustaceans**

Source : wiseoceans.com/IUCN

© AFP

Factfile on Cuvier's beaked whale, repeatedly stranded in the Mediterranean because of man-made sonar, according to a new theory

'Atypical' mass strandings

Developed in the 1950s to detect submarines, mid-frequency active sonar (MFAS) is used today in naval patrols and exercises, especially by

the United States and its NATO allies.

Starting around 1960, ships began emitting underwater signals in a range of about 5 kilohertz (kHz).

That is when the mass beaching of beaked whales, especially in the Mediterranean, began.

Between 1960 and 2004, 121 of these so-called "atypical" mass strandings took place, with at least 40 closely linked in time and place with naval activities.

These were not individual strandings of old or sick animals, nor en masse strandings such the one last November in New Zealand, when more than 200 pilot whales beached themselves together.

Rather, a handful or more [beaked whales](#) would wash ashore within a day or two, and no more than few dozen kilometres apart.

The most deadly episode, in 2002, saw 14 stranded over a 36-hour period in the Canary Islands during a NATO naval exercise.

"Within a few hours of the sonar being deployed, the animals started showing up on the beach," Bernaldo de Quiros said.

Outwardly, the whales showed no signs of disease or damage: they had normal body weight, and no skin lesions or infections.



Starting around 1960, ships began emitting underwater signals in a range of about 5 kilohertz (kHz) which is when the mass beaching of beaked whales began

Internally, it was another story. Nitrogen gas bubbles filled the veins, and their brains were ravaged by haemorrhaging.

Autopsies also revealed damage to other organs, as well as to the spinal cord and central nervous system.

Canary Island moratorium

As with altitude sickness, reactions—in humans, and probably in

whales—to nitrogen bubbles in the blood vary in type and intensity.

A 2003 study in *Nature* on the possible link between sonar and whale deaths led to Spain banning such naval exercises around the Canary Islands in 2004.

"Up until then, the Canaries were a hotspot for this kind of 'atypical' strandings," said Bernaldo de Quiros. "Since the moratorium, none have occurred."

The authors called for similar bans to be extended to other regions where at-risk [whales](#) are known to congregate.

The Cuvier's grows up to seven metres (23 feet) and dines mainly on deep-water squid and fish. Its upwardly turned mouth gives the impression of a permanent smile.

The whale is listed as "vulnerable" on the IUCN Red List of endangered species, and is thought to have a global population of 5,000 to 7,000.

Other threats include ship strikes, ocean pollution and shifting habitats caused by [climate change](#).

More information: Advances in research on the impacts of anti-submarine sonar on beaked whales, *Proceedings of the Royal Society B*, [rspb.royalsocietypublishing.org1098/rspb.2018.2533](https://royalsocietypublishing.org/doi/10.1098/rspb.2018.2533)

© 2019 AFP

Citation: For some whales, sonar may provoke suicidal behaviour: study (2019, January 30) retrieved 19 September 2024 from <https://phys.org/news/2019-01-whales-sonar-provoke-suicidal-behaviour.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.