


# These strange fossils are closely related to sea urchins

February 4 2019

---




Stunningly well-preserved fossilized soft tissues of a stylophoran have recently been discovered. Shown here is the reconstruction of an individual of the stylophoran genus *Thoralicystis*. Stylophorans measured 0.5 to 4 cm and had flat, massive bodies or tests with paddle-like extensions, analogous to snowshoes, which allowed them to stay over soft seafloors. Credit: Rich Mooi / California Academy of Science

Just a few centimeters long, these animals thrived in the ocean roughly

half a billion years ago. Because of their odd morphology, scientists have long struggled to find their branch on the tree of life.

Was their long appendage similar to a tail? That would make them ancestors of the vertebrates. However, their skeletons are made up of many calcite plates, suggestive of the bodies of echinoderms like [sea urchins](#) and starfish, even though they lack the characteristic symmetry of these animals.

A team led by Bertrand Lefebvre, a CNRS researcher at the Laboratoire de Géologie de Lyon, could finally settle this 150-year-old debate, using exceptionally preserved fossils from the Bou Izargane excavation in Morocco. Very unusually, the [soft tissues](#) of the fossilized creatures were preserved as pyrite, a ferrous mineral. By mapping the distribution of iron within the fossils, the researchers were able to clarify the fine structure of the appendage, which turns out to be comparable to that of a starfish arm. So these organisms had neither a head nor a tail, but rather a feeding arm.



Mapping of iron distribution in a portion of the articulated appendage of a stylophoran from the Moroccan region of Zagora. Iron-rich pyritized zones appear in green. They indicate where living soft tissue once laid. Here, we can clearly see ambulacral podia (pieds ambulacraires)—hollow tubes for feeding and locomotion—branching out along the water vascular canal (canal aquifère).

The arms of starfish and other echinoderms, like crinoids and brittle stars, possess the same type of structure. Credit: Bertrand Lefebvre / LGL-TPE / CNRS

**More information:** Bertrand Lefebvre et al. Exceptionally preserved soft parts in fossils from the Lower Ordovician of Morocco clarify stylophoran affinities within basal deuterostomes, *Geobios* (2018). [DOI: 10.1016/j.geobios.2018.11.001](https://doi.org/10.1016/j.geobios.2018.11.001)

Provided by CNRS

Citation: These strange fossils are closely related to sea urchins (2019, February 4) retrieved 16 April 2024 from <https://phys.org/news/2019-02-strange-fossils-sea-urchins.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.