

Revising the history of big, climate-altering volcanic eruptions

5 February 2019



Researchers drilled several ice cores from Dome C, one of the highest points on the Antarctic ice sheet, to reconstruct records of large volcanic eruptions stored in layers of ice. The researchers' new method is more accurate than previous techniques, providing a new perspective on the past 2,600 years of Earth's volcanic history. Credit: Bruno Jourdain

For all their destructive power, most volcanic eruptions are local events. Lava flows tend to reach only a few miles at most, while airborne ash and soot travel a little farther. But occasionally, larger eruptions can launch particles into the stratosphere, more than 6 miles above Earth's surface. The 1991 eruption of Mount Pinatubo in the Philippines—the world's largest eruption in the past 100 years—is a prime example of a stratospheric eruption.

When volcanic particles reach the stratosphere they stay aloft for a long time, reflecting sunlight and temporarily cooling the planet. By understanding the history of these big eruptions, researchers can begin to place short cooling episodes and other discrete climate events into the context of large-scale climate patterns.

Researchers working at the University of Maryland,

the Université Grenoble Alpes in France, the Ecole Normale Supérieure in France and the Tokyo Institute of Technology have devised a new, more accurate system for identifying large stratospheric eruptions recorded in the layers of Antarctic ice cores.

Using their method, the researchers made some important revisions to the known history of big eruptions—correcting the record on several misidentified events while discovering a few as yet unknown stratospheric eruptions. The researchers described their approach, which identifies airborne volcanic particles with a specific chemical signature, in a paper published January 28, 2019, in the journal *Nature Communications*.

"I find it very exciting that we are able to use chemical signals to build a highly accurate record of large, climate-relevant stratospheric eruptions," said James Farquhar, a professor of geology at UMD and a co-author of the research paper. "This historical record will be highly useful for climate scientists seeking to understand the role of large eruptions in climate oscillations. But there is also the basic marvel of reading a chemical fingerprint that is left behind in ice."

Eventually, volcanic particles fall from the stratosphere, settling on the ground below. When they land on snow, the particles get covered up by more snow that gets compacted into ice. This preserves a record of the [eruption](#) that survives until the ice melts. Researchers can drill and retrieve ice cores in places like Antarctica and Greenland, revealing eruption records that stretch back several thousand years.



This photo, taken on June 12, 1991, shows the eruption column of Mount Pinatubo on Luzon Island in the Philippines. The eruption--the largest on Earth in the past 100 years--ejected particles into the stratosphere, more than 6 miles above the planet's surface. New research uses ice core data to rewrite the past 2,600 years of large stratospheric eruptions like this one. Credit: Dave Harlow/USGS

Because particles from large stratospheric eruptions can spread across the globe before falling to the ground, previous methods identified stratospheric eruptions by looking for sulfate particle layers in ice from both hemispheres—usually from Antarctica and Greenland. If the same layers of sulfate showed up in both cores, deposited at the same time in Earth's history, researchers would conclude that the particles came from the same large, stratospheric eruption.

"For eruptions that are intense enough to inject material into the stratosphere, there is a telltale signature in the sulfur isotope ratios of sulfate preserved in ancient ice layers," explained Farquhar, who also has an appointment in UMD's Earth System Science Interdisciplinary Center. "By instead focusing on this distinct sulfur isotope signature, our new method yielded some surprising and useful results. We found that prior reconstructions missed some stratospheric events and falsely identified others."

The study's lead author, Elsa Gautier from the Université Grenoble Alpes, did a significant portion

of the analyses at UMD while on a Fulbright scholarship to work with Farquhar in 2013. Following Gautier's lead, the researchers developed their method using ice cores collected at a remote site in Antarctica called Dome C. One of the highest points on the Antarctic ice sheet, Dome C is home to ice layers that stretch back nearly 50,000 years.

Gautier and her colleague Joel Savarino, also at the Université Grenoble Alpes, collected ice cores at Dome C that contain records stretching back roughly 2,600 years, covering a large portion of recorded human history.

The researchers used their method to confirm that many events had indeed been properly identified by the older method of matching up corresponding sulfate layers in ice cores from both hemispheres. But some events, formerly thought to be big stratospheric eruptions, did not have the telltale sulfur isotope signature in their sulfate layers. Instead, the researchers concluded, these layers must have been deposited by two or more smaller volcanoes that erupted at about the same time at high latitudes in both hemispheres.

The researchers also found some big stratospheric events that contain the isotope signature, but were somehow constrained to the Southern Hemisphere.

"This is a strength of our approach, because these events would have a climate impact but are missed by other methods," Farquhar said. "We have made a significant improvement to the reconstruction of large stratospheric eruptions that occurred over the past 2,600 years. This is critically important for understanding the role of [volcanic eruptions](#) on climate and possibly for understanding certain events in human history, such as widespread famines. It can also help to inform future climate models that will take large volcanic events into account."

The research paper, "2600 years of stratospheric volcanism through sulfate isotopes," Elsa Gautier, Joel Savarino, Joost Hoek, Joseph Erbland, Nicolas Caillon, Shohei Hattori, Naohiro Yoshida, Emanuelle Albalat, Francis Albarede and James Farquhar, was published in the journal

Nature Communications on January 28, 2019.

More information: E. Gautier et al, 2600-years of stratospheric volcanism through sulfate isotopes, *Nature Communications* (2019). [DOI: 10.1038/s41467-019-08357-0](#)

Provided by University of Maryland
APA citation: Revising the history of big, climate-altering volcanic eruptions (2019, February 5) retrieved 27 May 2019 from <https://phys.org/news/2019-02-history-big-climate-altering-volcanic-eruptions.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.