

Scientists advance new technology to protect drinking water from Lake Erie algal toxins

11 February 2019



Dr. Jason Huntley, associate professor in The University of Toledo Department of Medical Microbiology and Immunology in the UT College of Medicine and Life Sciences. Credit: Dan Miller, The University of Toledo

Before the 2014 Toledo Water Crisis left half a million residents without safe drinking water for three days, Dr. Jason Huntley's research at The University of Toledo focused on bacteria that cause pneumonia.

After the harmful algal bloom prompted the city of Toledo's "Do Not Drink" advisory, the microbiologist expanded his [research projects](#) to target microcystin.

"I live here, and I have a young son," said Huntley, associate professor in the UT Department of Medical Microbiology and Immunology in the College of Medicine and Life Sciences. "I don't want toxins in the [water](#), and I am committed to helping the [water treatment plant](#) protect the public."

Huntley's research lab recently made major

progress in his mission to create a biofilter that uses naturally occurring Lake Erie bacteria to remove microcystin released by [harmful algal blooms](#) from drinking water, reducing or eliminating the use of chlorine and other chemicals.

"We've identified groups of bacteria in Lake Erie that can be used to naturally purify water. To our knowledge, these bacteria have not been previously used to fight harmful algal blooms in other parts of the world," Huntley said.

The microbiologists successfully isolated bacteria from Lake Erie that degrade the microcystin toxin known as MC-LR—the most toxic, most common and most closely linked to liver cancer and other diseases—at a daily rate of up to 19 parts per billion (ppb).

Water analysts and toxicologists measure microcystin and other contaminants using the metric of ppb; one ppb is one part in 1 billion. These ppb numbers are important for [human health](#) because the U.S. Environmental Protection Agency recommends that young children not drink water containing more than 0.3 ppb of microcystin and adults not drink water containing more than 1.6 ppb of microcystin.

"The bacteria we've identified can degrade much more toxin than was reported in the 2014 water crisis," Huntley said. "Based on recorded toxin levels in Lake Erie in recent years, these rates would be able to effectively remove microcystin from water supplies."

None of the 13 microcystin-degrading bacterial isolates has been associated with human disease, so their use in future water-purifying biofilters is unlikely to be a public health concern. The identified bacteria include *Flectobacillus major*,

Pseudomonas lutea, *Agrobacterium albertimagni*, *Leadbetterella byssophila*, *Pseudomonas putida*, *Flectobacillus major*, *Pseudomonas hunanensis*, *Runella slithyformis*, *Porphyrobacter sp.*, *Pseudomonas parafulva*, *Sphingobium yanoikuyae*, *Pseudomonas fluorescens* and *Sphingobium yanoikuyae*.

however, Huntley said those specific types of bacteria were not found in any of his Lake Erie studies.

Thirteen water samples used for the study were collected from visible algal blooms in the summers of 2014 and 2015 in the western basin of Lake Erie. The scientists added MC-LR to each water sample every three-to-four days for approximately four weeks, alongside a control group that did not receive additional MC-LR.

The lab used multiple approaches to confirm the microcystin degradation results, including mass spectrometry and the ELISA test, which is the standard method water treatment plant operators use to measure microcystin concentration during algal bloom season.

His lab is now in the process of identifying the enzymatic pathways the bacteria use to break down microcystin.

Currently, municipal water treatment plants remove or degrade microcystin using methods such as chlorination, ozonation, activated carbon adsorption and flocculation.

"Those techniques are not ideal because of high costs, limited removal efficiencies, and they lead to the production of harmful byproducts or hazardous waste," Huntley said. "Biofilters are a cost-effective and safe alternative to the use of chemicals and other conventional water treatment practices."

"We're very excited about the research and the findings," said Andrew McClure, administrator for the city of Toledo's Collins Park Water Treatment Plant. "We've had preliminary talks with Dr. Huntley about ways we can implement it as a treatment technique in our plant's process."

Huntley's team is developing and testing biofilters—water filters containing the specialized [bacteria](#) that degrade microcystin toxins from lake water as it flows through the filter. Huntley holds a provisional patent on this technology.

The research was supported by grants from the Ohio Department of Higher Education through the



The lab-scale biofilters used during Dr. Jason Huntley's research at The University of Toledo are sand filters that contain biologically active bacteria that break down microcystin toxins. Credit: Dr. Jason Huntley, The University of Toledo

The research is published in the February issue of the *Journal of Great Lakes Research*.

Researchers in Australia, China and other countries also have identified bacteria that can chew up and break down microcystin from algal blooms,

state's Harmful Algal Bloom Research Initiative, which consists of 54 science teams at universities across the state seeking solutions to address toxic algae in Lake Erie.

Provided by University of Toledo

APA citation: Scientists advance new technology to protect drinking water from Lake Erie algal toxins (2019, February 11) retrieved 20 November 2019 from <https://phys.org/news/2019-02-scientists-advance-technology-lake-erie.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.