

# New deep sea animal discoveries warrant expanded protections in Costa Rican waters

11 February 2019



Beautiful school of Cutlassfish, vertically feeding in the water column. Lit up by the lights on ROV SuBastian while exploring on the #CostaRicaDeep expedition. Credit: Schmidt Ocean Institute

A three week expedition off the coast of Costa Rica has just expanded our knowledge of deep sea ecosystems in the region. Led by Dr. Erik Cordes, Temple University, the scientists aboard research vessel Falkor surveyed the continental margin for seamounts and natural gas seeps, where specialized biological communities are found. The seamounts extending from the mainland to the Cocos Islands National Park provide an important corridor for the animals occupying the area.

Investigating these systems on all biological size scales, the team focused on relationships between [species](#), from microbes to fauna like fish and corals. At least four new species of deep-sea corals and six other animals that are new to science were found. This expedition represents the first time that seven of the seamounts in the area have been surveyed. The [survey results](#), including description of the coral communities that they host, will support the effort to create a new marine protected area around these seamounts ensuring that they are not impacted by fishing or potential mining activities.

"This research will support Costa Rica's efforts to conserve these important habitats by providing a baseline of the incredible species and ecosystems found in the deeper areas that don't always attract the attention that they deserve," said Schmidt Ocean Institute Cofounder Wendy Schmidt. "One of the most important things we can do now is understand how these communities work, so if there are changes in the future we can measure human impact."



Chimaeras are cartilaginous fish, largely confined to deep water. Their closest living relatives are sharks, though their last common ancestor with sharks lived nearly 400 million years ago. Credit: Schmidt Ocean Institute

Even in [deep waters](#), humans pose a threat to these fragile ecosystems. During one of the 19 remotely operated vehicle dives the accumulation of trash at 3,600 meters depth (more than 2 miles) was discovered. Threats to the deep sea already exist, including fishing and energy industries that are moving into deeper water, and the persistent risk of climate change. There are rare organisms and spectacular habitats on the seamounts; it is important to preserve them before they are impacted by these and other threats.

One unique discovery during the expedition was

the consistent zonation of seamounts related to the amount of oxygen present. Decreasing oxygen in the ocean due to a warming planet may eventually affect these zones dominated by corals, sea fans, sponges, brittle stars and small oysters. "Every dive continues to amaze us," said Cordes. "We discovered species of reef-building stony corals at over 800 meters depth on two different seamounts. The closest records of this species are from the deep waters around the Galapagos Islands. The deep sea is the largest habitat on Earth. Understanding how that habitat functions will help us to understand how the planet as a whole works."

Provided by Schmidt Ocean Institute

APA citation: New deep sea animal discoveries warrant expanded protections in Costa Rican waters (2019, February 11) retrieved 26 September 2021 from <https://phys.org/news/2019-02-deep-sea-animal-discoveries-warrant.html>

*This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.*