


Total synthesis of marine antitumor agents trabectedin and lurbinectedin

19 February 2019



Credit: Wiley

With its vast numbers of different lifeforms, the sea is a largely unexplored source of natural products that could be starting points for new pharmaceuticals, such as the antitumor drugs trabectedin and lurbinectedin. Because only tiny amounts can be obtained from sea organisms, synthetic production is necessary. In the journal *Angewandte Chemie*, scientists have introduced a new, efficient synthetic route for these two drugs. A key step is the light-controlled activation of a carbon–hydrogen bond.

Trabectedin (also called ecteinascidin) comes from the sea squirt species *Ecteinascidia turbinata* and is the first marine natural product to be used clinically as a [drug](#)—for the treatment of advanced soft tissue sarcoma. Lurbinectedin has a slightly modified structure and is currently in phase III clinical studies for the treatment of certain lung and breast cancers. One ton of [sea squirts](#) are needed to acquire about one gram of trabectedin. A viable and efficient synthetic route for making this and related drugs in sufficient quantities is therefore needed urgently. However, trabectedin has thus far proven to be one of the most challenging target molecules in natural products synthesis. Various synthetic routes have been proposed but none is

really viable. Current methods are very complex, require expensive and uncommon reagents, and deliver unsatisfactory yields.

Researchers working with Dawei Ma at the Shanghai Institute of Organic Chemistry (China) have now described a more efficient and viable de novo synthetic route for trabectedin and lurbinectedin. De novo, also known as [total synthesis](#), means that the natural product is completely synthesized from small, common starting materials.

The synthesis starts with the amino acid S-tyrosine and consists of 26 individual steps. First, several steps are used to produce an intermediate, which acts as the starting material for the separate production of the two halves of the target molecule—Trabectedin or Lurbinectedin. These are then bound together in a later reaction step.

The key step of the [synthesis](#) is the light-controlled activation of a normally unreactive carbon–hydrogen bond (remote C–H activation). A radical rearrangement mechanism leads to a ring closure in which a quinone group is converted into a 1,3-benzodioxole unit, which is a structural component found in many natural products. The reaction was particularly efficient under irradiation with [blue light](#) in tetrahydrofuran solvent.

The scientists hope that their synthetic route offers a practical and economical method for the production of trabectedin and lurbinectedin, finally providing adequate supplies of these complex marine antitumor drugs.

More information: Weiming He et al. A Scalable Total Synthesis of the Antitumor Agents Et-743 and Lurbinectedin, *Angewandte Chemie International Edition* (2019). DOI: [10.1002/anie.201900035](https://doi.org/10.1002/anie.201900035)

Provided by Wiley

APA citation: Total synthesis of marine antitumor agents trabectedin and lurbinectedin (2019, February 19) retrieved 13 May 2021 from <https://phys.org/news/2019-02-total-synthesis-marine-antitumor-agents.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.