

# Old for new, using ancient genetic variation to supercharge wheat

March 27 2019

---



Credit: CC0 Public Domain

Wheat is responsible for half of global calories consumed either directly or as animal feed and we need to make a lot more of it in the coming decades. Now, an important paper marks a step change in how breeders

might approach wheat breeding using previously untapped sources of variation to increase yields to meet demand.

The recent paper, "Elucidating the genetic basis of biomass accumulation and radiation use efficiency in spring wheat and its role in yield potential", first-authored by Dr. Ryan Joynson of EI's Anthony Hall Group and Dr. Gemma Molero of Matthew Reynold's team at CIMMYT, presents a valuable step forward for researchers interested in increasing wheat yield potential: a crucial task if we are to sustainably feed the planet.

The research, funded through IWYP (International Wheat Yield Partnership) shines light on the role of radiation use efficiency (RUE, how sunlight is converted into plant mass) in increasing the yield potential of wheat: how we can increase plant biomass without sacrificing grain yield—a major challenge.

This research highlights:

- Identification of genomic regions affecting RUE and biomass accumulation in spring wheat
- How increases in RUE can be achieved through strategic integration of 'exotic' wheat into breeding programmes
- Identification of common genomic regions affecting yield, biomass and radiation use efficiency
- Enrichment of genes related with photoprotection in genomic regions associated with RUE at various growth stages
- Identification of untapped variation in wheat that can be harnessed and used by scientists and breeders

Dr. Ryan Joynson, co-first author on the paper, said: "It's really gratifying to use my computational skills in such an applied project and to see this research being incorporated into important international

breeding programmes."

## **The challenge**

Thanks to Norman Borlaug's green revolution, led from CIMMYT, the development and improvement of dwarf varieties of wheat have led to a drastic increase in wheat yields over the better half of the 21st century. Yet, this consistent increase has wavered in recent years.

Continually increasing wheat yields is challenging, increasingly so when we must more-than-double them if we are to sustainably feed 10 billion people by 2050, half of whom will rely on wheat as a staple for bread, noodles and more. At current rates, we are predicted to only get to 38% increased wheat yields by 2050, which would mean a significant shortfall.

The challenge is the well-known trade-off between things such as grain weight/number and biomass. How do we reduce the leafy portion of a plant, or the underground portion, enough to boost harvestable product such that other aspects, including radiation use efficiency (hence photosynthesis) and nutrient uptake, are not disrupted or diminished?

Alternatively, how do we increase biomass and the relative efficiency of photosynthesis, without reducing the harvestable portion of the crop—the grain?

Photosynthesis itself is pretty inefficient, so it must be possible to find a happy medium between less biomass, more grain, and more efficient photosynthesis—so where better to start than to identify genes and genetic regions underpinning all of these traits in elite lines of wheat?

By growing 150 types of wheat, then mapping the differences in growth to differences at the genetic level, it was possible to understand areas of

commonality in desired traits for crop improvement.

## **A new resource for breeders**

Through analysing new variation in elite wheat lines, while looking for the genetic markers underpinning and linking traits such as radiation use efficiency, biomass, yield, grain number, grain weight, etc., the researchers have provided an important resource for scientists and breeders to exploit.

Namely, markers for use in marker-assisted breeding, that will help increase desired outcomes—such as biomass, thousand grain weight and radiation use efficiency—while avoiding the trade-off between grain weight and grain number, or between biomass and harvest index.

The study represents the completion of two years of work looking at 150 wheat varieties, using a genome-wide association study to analyse 31 different traits. Among the genetic regions identified, several were found in specific regions common to grain yield, biomass and radiation use efficiency.

Another important aspect of the work was the investigation of the effect of sources of new variation into elite lines of wheat. Comparing elite wheat varieties alongside 'exotic' lines such as landraces and synthetic wheat, the researchers pinpointed areas of interest for future breeding programmes, i.e. where we can introduce more variation into our best [wheat](#) varieties for desired outcomes.

**More information:** Gemma Molero et al. Elucidating the genetic basis of biomass accumulation and radiation use efficiency in spring wheat and its role in yield potential, *Plant Biotechnology Journal* (2018). [DOI: 10.1111/pbi.13052](https://doi.org/10.1111/pbi.13052)

Provided by Earlham Institute

Citation: Old for new, using ancient genetic variation to supercharge wheat (2019, March 27)  
retrieved 19 September 2024 from

<https://phys.org/news/2019-03-ancient-genetic-variation-supercharge-wheat.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.