


Researchers remove harmful hormones from Las Vegas wastewater using green algae

April 8 2019



Xuelian Bai, Ph.D., Assistant Research Professor of Environmental Sciences, works with an algae sample in the Environmental Engineering Laboratory at the Desert Research Institute in Las Vegas. Credit: Sachiko Sueki.

A common species of freshwater green algae is capable of removing

certain endocrine disrupting chemicals (EDCs) from wastewater, according to new research from the Desert Research Institute (DRI) in Las Vegas.

EDCs are natural hormones and can also be found in many plastics and pharmaceuticals. They are known to be harmful to wildlife, and to humans in large concentrations, resulting in [negative health effects](#) such as lowered fertility and increased incidence of certain cancers. They have been found in trace amounts (parts per trillion to parts per billion) in treated wastewater, and also have been detected in [water samples](#) collected from Lake Mead.

In a new study published in the journal *Environmental Pollution*, DRI researchers Xuelian Bai, Ph.D., and Kumud Acharya, Ph.D., explore the potential for use of a species of freshwater green [algae](#) called *Nannochloris* to remove EDCs from treated wastewater.

"This type of algae is very commonly found in any freshwater ecosystem around the world, but its potential for use in wastewater treatment hadn't been studied extensively," explained Bai, lead author and Assistant Research Professor of environmental sciences with the Division of Hydrologic Sciences at DRI. "We wanted to explore whether this species might be a good candidate for use in an algal pond or constructed wetland to help remove wastewater contaminants."



Samples of Nannochloris grow in the Environmental Engineering Laboratory at the Desert Research Institute in Las Vegas. This species of green algae was found to be capable of removing certain types of endocrine disrupting chemicals from treated wastewater. Credit: Xuelian Bai/DRI

During a seven-day laboratory experiment, the researchers grew Nannochloris algal cultures in two types of treated wastewater effluents collected from the Clark County Water Reclamation District in Las Vegas, and measured changes in the concentration of seven common EDCs.

In wastewater samples that had been treated using an ultrafiltration

technique, the researchers found that the algae grew rapidly and significantly improved the removal rate of three EDCs (17 β -estradiol, 17 α -ethinylestradiol and salicylic acid), with approximately 60 percent of each contaminant removed over the course of seven days. In wastewater that had been treated using ozonation, the algae did not grow as well and had no significant impact on EDC concentrations.

One of the EDCs examined in the study, triclosan, disappeared completely from the ultrafiltration water after seven days, and only 38 percent remained in the ozonation water after seven days—but this happened regardless of the presence of algae, and was attributed to breakdown by photolysis (exposure to light).

"Use of algae for removing heavy metals and other inorganic contaminants have been extensively studied in the past, but for removing organic pollutants has just started," said Acharya, Interim Vice President for Research and Executive Director of Hydrologic Sciences at DRI.

"Our research shows both some of the potential and also some of the limitations for using *Nannochloris* to remove EDCs from wastewater."



Researcher examines a sample of quagga mussels collected from Lake Mead. A recent study by Bai and Acharya found that endocrine disrupting chemicals are accumulating in the body tissues of these mussels. Credit: Xuelian Bai/DRI

Although these tests took place under laboratory conditions, a previous study by Bai and Acharya that published in November 2018 in the journal *Environmental Science and Pollution Research* examined the impacts of these same seven EDCs on quagga mussels (*Dreissena bugensis*) collected from Lake Mead. Their results showed that several of the EDCs (testosterone, bisphenol A, triclosan, and [salicylic acid](#)) were accumulating in the body tissues of the mussels.

"Algae sit at the base of the food web, thereby providing food for organisms in higher trophic levels such as [quagga mussels](#) and other zooplanktons," Bai said. "Our study clearly shows that there is potential for these contaminants to biomagnify, or build up at higher levels of the food chain in the aquatic ecosystem."

Bai is now working on a new study looking for antibiotic resistance in genes collected from the Las Vegas Wash, as well as a study of microplastics in the Las Vegas Wash and Lake Mead. Although Las Vegas's treated wastewater meets Clean Water Act standards, Bai hopes that her research will draw public attention to the fact that treated wastewater is not 100 percent clean, and will also be helpful to utility managers as they develop new ways to remove untreated contaminants from wastewater prior to release.

"Most wastewater treatment plants are not designed to remove these unregulated contaminants in lower concentrations, but we know they may cause health effects to aquatic species and even humans, in large concentrations," Bai said. "This is concerning in places where [wastewater](#) is recycled for use in agriculture or released back into drinking water sources."

More information: Xuelian Bai et al, Removal of seven endocrine disrupting chemicals (EDCs) from municipal wastewater effluents by a freshwater green alga, *Environmental Pollution* (2019). [DOI: 10.1016/j.envpol.2019.01.075](#)

Provided by Desert Research Institute

Citation: Researchers remove harmful hormones from Las Vegas wastewater using green algae (2019, April 8) retrieved 19 September 2024 from <https://phys.org/news/2019-04-hormones-las->

vegas-wastewater-green.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.