

Keeping the taste, reducing the salt

April 11 2019, by Scott Weybright



While humans need the salt in snacks like potato chips, Americans consume significantly more salt than is necessary or even healthy. Credit: Public Domain

Washington State University researchers have found a way to make food taste salty but with less of the sodium chloride tied to poor health.

"It's a stealth approach, not like buying the 'reduced [salt](#)' option, which people generally don't like," said Carolyn Ross, a Food Science professor at WSU. "If we can stair-step people down, then we increase health while still making food that people want to eat."

In a paper published in the *Journal of Food Science*, Ross and colleagues looked at salt blends that use less sodium chloride and include other salts like calcium chloride and [potassium chloride](#).

Both of those salts have no [adverse health effects](#) on people, Ross said. Potassium can actually help reduce blood pressure. Unfortunately, they aren't very tasty.

"Potassium chloride, especially, tastes really bitter and people really don't like it," Ross said.

The researchers used tasting panels and WSU's electronic tongue to see just how much they could add of the replacement salts for standard sodium chloride before people found the food unacceptable to eat.

Some tasting panels tested a variety of salt solutions, or salt in water, while others tested different salt combinations in tomato soup.



Carolyn Ross Credit: WSU

Using the e-tongue and panels, they found that a blend using approximately 96.4 percent sodium chloride with 1.6 percent potassium chloride and 2 percent calcium chloride was the ideal reduction.

They had a higher reduction when they added only calcium chloride, getting acceptable rates with a combination of 78 percent sodium chloride and 22 percent [calcium chloride](#).

"This combination of the two salts did not significantly differ compared

to 100 percent [sodium chloride](#)," Ross said. "But when we added potassium chloride, consumer acceptance decreased."

While humans need salt, Americans consume significantly more than is necessary or even healthy. According to the U.S. Office of Disease Prevention and Health Promotion, the recommended maximum amount of salt consumed per day is less than 2,300 mg. The average American adult female consumes 2,980 mg per day, while males average over 4,000 mg per day.

Recent findings have suggested that gradual reductions in salt over a period of years is the best way to reduce salt consumption. Using one of the new blends for a specified time frame could lead to greater reductions down the road.

More information: Sasha M. Barnett et al, Identification of a Salt Blend: Application of the Electronic Tongue, Consumer Evaluation, and Mixture Design Methodology, *Journal of Food Science* (2019). [DOI: 10.1111/1750-3841.14440](#)

Provided by Washington State University

Citation: Keeping the taste, reducing the salt (2019, April 11) retrieved 20 September 2024 from <https://phys.org/news/2019-04-salt.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.