

# Worldwide study reveals air pollution link to unborn baby growth

18 April 2019



Credit: CC0 Public Domain

Exposure to air pollution is linked to babies' growth during pregnancy, according to a landmark new study.

Scientists at the University of Aberdeen have reviewed over a decade's worth of [research evidence](#) from around the world to establish the extent to which mothers' exposures to [air pollution](#), diet, alcohol and chemicals affects fetal growth.

While there was [insufficient evidence](#) that poor diet, alcohol or chemicals were factors, the study found evidence that exposure to nitrogen dioxide resulted in smaller fetal head size, particularly in the last three months of pregnancy.

Nitrogen dioxide is primarily generated by vehicle traffic, but can be present in the home from cigarette smoke, or butane and kerosene heaters and stoves.

Professor Steve Turner, who led the study, said: "In our research we looked at all the studies that measured the effects of mothers' exposures to everyday substances including air pollutants, alcohol, and diet on the size of the unborn baby, measured through ultrasound from half way through the pregnancy onwards.

"What was unique about our review of the literature is that we looked at unborn babies to see if mothers' exposures to these factors affected fetal development. We used [medical literature](#) dating back 13 years, when the first studies linking exposure to fetal measurements was first published.

"The seven studies where air pollution was measured and linked to fetal size were from different geographical areas of the world, including Australia, the USA and several countries in Europe. However in all of the studies the evidence was clear that in the third trimester in particular, exposure to [nitrogen](#) dioxide reduced fetal growth."

Professor Turner continued: "Previous research has shown that being small (for gestational age) at birth is associated with increased risk for conditions that include coronary artery disease, type II diabetes and asthma.

"Our research has shown that the link between [exposure](#) and fetal growth is apparent well before birth, so any potential interventions need to happen

in the early stages of pregnancy.

"Furthermore, the findings also suggest that public health measures are urgently required to minimise pregnant mothers' exposures to [nitrogen dioxide](#)."

**More information:** Ivory Huang et al. A systematic review of associations between maternal exposures during pregnancy other than smoking and antenatal fetal measurements, *Environmental Research* (2019). [DOI: 10.1016/j.envres.2019.04.005](#)

Provided by University of Aberdeen

APA citation: Worldwide study reveals air pollution link to unborn baby growth (2019, April 18) retrieved 19 September 2020 from <https://phys.org/news/2019-04-worldwide-reveals-air-pollution-link.html>

*This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.*