


Reducing energy required to convert CO₂ waste into valuable resources

April 22 2019



Chemical and biomolecular engineering professor and department chair Paul Kenis, right, and graduate student Shawn Lu are co-authors of a new study that examines the feasibility of a new CO₂ waste-to-value technology. Credit: L. Brian Stauffer

Surplus industrial carbon dioxide creates an opportunity to convert waste

into a valuable commodity. Excess CO₂ can be a feedstock for chemicals typically derived from fossil fuels, but the process is energy-intensive and expensive. University of Illinois chemical engineers have assessed the technical and economic feasibility of a new electrolysis technology that uses a cheap biofuel byproduct to reduce the energy consumption of the waste-to-value process by 53 percent.

The new findings are published in the journal *Nature Energy*.

Conversion of CO₂ to chemicals like ethylene for plastics is possible through a process called electrochemical reduction. Typically, a stream of CO₂ gas and a fluid electrolyte move through an electrolysis cell that breaks the CO₂ down into molecules like ethylene on the cathode, but it also produces oxygen from water on the anode, the researchers said.

"About 90 percent of the energy required in conventional CO₂ reduction is used up by the oxygen-producing, anode side of an electrolysis cell," said Paul Kenis, a chemical and biomolecular engineering professor, department chair and study co-author. "But there is no big market for the excess oxygen, so 90 percent of the energy is essentially wasted."

Finding a feed material that reduces the energy to drive the anode reaction could be a strategy for radically reducing the energy requirements of CO₂ conversion, according to a recent National Academies Report of which Kenis was a co-author.

The new study proposes glycerol—an organic byproduct of sugar cane biofuel production that requires less energy to oxidize—as an alternative to the energy-intensive oxygen-producing step.

To test if the new electrolysis technique has the potential to push the full CO₂ conversion process to a carbon neutral or negative budget, the researchers examined the cost and energy consumption for the

production cycle of the waste-to-value process. The four-step cycle includes the capture of industrial CO₂ waste gas, the input of electricity, the new

"Our model uses the current electrical grid setup as the source of electricity to make the scenario more realistic," Kenis said. "Being able to drive CO₂ conversion with already-in-place infrastructure—and not relying on the hope of the future grid being powered by 100 percent renewables—while achieving carbon neutrality or negativity could be a holy grail scenario."

The analysis includes best- and worst-case CO₂ emissions and energy consumption scenarios and concludes that the prospects of CO₂ reduction, in terms of CO₂ emissions and economics, can drastically improve by looking beyond conventional anode reactions.

"The glycerol-based electrolysis reaction shows a lot of promise. However, we will continue to explore other organic waste materials because even when production rises in the wake of increased biofuel production, it still will not be enough to fully support the need," Kenis said. "The good news is that the chemistry involved is flexible and there are a lot of organic waste products that can do the job."

Many researchers focus on improving the selectivity and activity of chemical catalysts for CO₂ reduction reactions, and that work needs to continue, said Sumit Verma, a former chemical and biomolecular engineering graduate student and study co-author. "Looking beyond oxygen evolution at the anode seems like a win-win situation, as we not only reduce the processes' [energy](#) consumption but also produce a second valuable product stream," he said.

More information: Co-electrolysis of CO₂ and glycerol as a pathway to carbon chemicals with improved technoeconomics due to low

electricity consumption, *Nature Energy* (2019). DOI: [10.1038/s41560-019-0374-6](https://doi.org/10.1038/s41560-019-0374-6),
www.nature.com/articles/s41560-019-0374-6

Provided by University of Illinois at Urbana-Champaign

Citation: Reducing energy required to convert CO₂ waste into valuable resources (2019, April 22) retrieved 25 April 2024 from <https://phys.org/news/2019-04-energy-required-co2-valuable-resources.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.