

Researchers trace 3,000 years of monsoons through shell fossils

April 26 2019



Ostracod shells from juvenile to adult used for analysis. Ostracod is an arthropod that has lived for 500 million years. (photo by Kentaro Kuroki of Yamada lab.)
Credit: Katsura Yamada, Shinshu University, Japan

The tiny shells at the bottom of Lake Nakaumi in southwest Japan may contain the secrets of the East Asia Summer Monsoon. This rainy season is fairly predictable, ushering in air and precipitation conducive to growing crops, but—sometimes without any hint—the pattern fails. Some areas of East Asia are left without rainfall, and their crops die. Other areas are inundated with rain, and their crops and homes flood.

Ostracoda shells are smaller than the white wisp a fingernail grows over a month, yet they have recorded the effects of sunshine and climate shifts for almost 500 million years.

A team of researchers dug into the lake and the rich historical record in the shells to better understand why East Asian summer monsoons vary at the centennial scale, which should hold relatively steady. They published their results on March 22, 2019, in *Scientific Reports*.

"The mechanisms driving the variations in East Asian summer monsoons remain unclear, so we used the [oxygen isotopes](#) from adult ostracode shells to reconstruct the variations over the last 3,000 years in southwestern Japan," said Katsura Yamada, paper author and a professor in the department of geology and faculty of science at Shinshu University.

Yamada and the team cored sections of the lake, retrieving shells from present day to 3,000 years ago. The scientists analyzed the shells, measuring a specific ratio between slightly different versions of oxygen, called isotopes. The isotope ratio can offer a glimpse into the atmosphere's precise composition thousands of years ago. A higher rate of nitrogen in the atmosphere will produce a different isotope of oxygen than times when nitrogen is less abundant.




Ostracod shells were collected from the bottom of the lake Nakaumi. Credit: Katsura Yamada, Shinshu University, Japan

The researchers found that the primary factor of the centennial-scale variations in the East Asia summer monsoon was solar activity, also called insolation.

"Our results and compiled data propose that insolation variation was a primary factor of the centennial-scale East Asia summer monsoon variations," Yamada said. "However, dominant factors affecting the variations can shift according to the solar insolation decreases."

During sunny periods, the insolation dominates the East Asian monsoon pattern. During cooling off periods, usually around glacial ice ages, other factors—such as wind patterns—took over as the dominant influencer.

"Our next goal is to clarify the relationship between East Asian [monsoon](#) variations and other climatic phenomena," Yamada said.



Ostracod shell fossils. Credit: Katsura Yamada, Shinshu University, Japan

This work was supported by the Japan Society for the Promotion of Science and the cooperative research program for the Center for Advanced Marine Core Research at Kochi University.

Other authors include Kazuma Kohara of the department of geology and Faculty of Science at Shinshu University; Minoru Ikehara of the Center for Advanced Marine Core Research at Kochi University; and Koji Seto of the Estuary Research Center at Shimane University.

More information: Katsura Yamada et al, The variations in the East Asian summer monsoon over the past 3 kyrs and the controlling factors, *Scientific Reports* (2019). [DOI: 10.1038/s41598-019-41359-y](https://doi.org/10.1038/s41598-019-41359-y)

Provided by Shinshu University

Citation: Researchers trace 3,000 years of monsoons through shell fossils (2019, April 26) retrieved 26 April 2024 from <https://phys.org/news/2019-04-years-monsoons-shell-fossils.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.