

3-D technology looks into the distant past

20 May 2019



Mesopotamian barb (*Capoeta damascina*) from Lake Homs, Syria. Scale shown = 1cm. Credit: Ayvazyan et al., 2018. *J Zool Syst Evol Res*

Researchers from the University of Tübingen and their colleagues from Switzerland have studied hundreds of fossil carp teeth for the first time using 3-D technologies. In 4 million-year old lake sediments from what is now the Armenian highlands, they found evidence of an astonishing variety of carp. Thanks to "virtual palaeontology," the researchers identified four closely-related species of Mesopotamian barb fish, so-called scrapers. They believe that these species lived together in a giant system of lakes and formed a "species flock:" a group of closely related species inhabiting the same biotope. It is the first time that a species flock of freshwater fish has been found in western Asia. Today, these carp species live in separate regions. Findings such as this can help us to understand evolutionary developments and the origins of species diversity.

The project was conducted by Anna Ayvazyan and Professor Madelaine Böhme of the Senckenberg Center for Human Evolution and Palaeoenvironment (HEP) at the University of

Tübingen, in cooperation with Dr. Davit Vasilyan of the Jurassica Museum in Porrentruy (Switzerland). The results of their study have been published in the latest *PLoS ONE*.

The scrapers (genus *Capoeta*) are an ecologically specialized family of carp. They have a specialized diet, feeding only on algae, which they scrape off stones. They live in rivers which flow through arid landscapes. Today, more than 30 species of scrapers are known. Until now, this variety of species was hard to explain; before the advent of 3-D analysis, it was impossible to identify fossil carp species.

Three-dimensional models were created with the help of computer tomography; the researchers were then able to investigate the structure and shape of the fossil carp teeth. The 3-D technology also provides high-resolution details of the tooth structures of living species. "This method enables us to identify fossil species for the first time," says the first author of the study, doctoral researcher Anna Ayvazyan. "We've only just discovered the amazing diversity of species that existed 4 million years ago."

The species diverged in the Caucasus

With this precise species identification, evolutionary models for fish faunas of the Middle East and the Caucasus can now be created for the first time. The researchers think that that the evolution of the scrapers is a result of complex interactions between geological, biological and environmental processes. These factors determine a region's species formation processes and the geographical distribution of species over several million years.

The researchers say that, over 5 million years ago in western Asia, a species complex of scrapers developed in a huge lake system. Due to the collision of the Arabian continental plate with Eurasia, what is now the Armenian highlands began to rise by several kilometers. Today it comprises eastern Anatolia, northwestern Iran,

Armenia, southern Georgia and western Azerbaijan, and is between 1,500 and 5,000 meters above sea level. These mountain formation processes divided the giant lake system, and the separated scraper species survived in the western Asian rivers flowing from the highlands: the Euphrates, Tigris, Kura and Aras. The [geological processes](#) led to the further differentiation and development of new species of Mesopotamian barb.

Biodiversity is threatened by human activity around the world. The recently-published UN Biodiversity Report lists one in three of more than 15,000 known freshwater fish species as endangered; most of them due to pollution, [climate change](#), overfishing and dam projects. This is particularly true of carp, the biggest family of freshwater fish worldwide. They occur only in very limited areas (endemism) and their distribution depends on being able to swim from one body of water to another. To prevent the extinction of the [species](#), it is important to understand which evolutionary mechanisms led to the emergence of this biodiversity.

More information: Anna Ayvazyan et al. Possible species-flock scenario for the evolution of the cyprinid genus *Capoeta* (Cypriniformes: Cyprinidae) within late Neogene lake systems of the Armenian Highland, *PLOS ONE* (2019). [DOI: 10.1371/journal.pone.0215543](#)

Provided by Universitaet Tübingen

APA citation: 3-D technology looks into the distant past (2019, May 20) retrieved 4 March 2021 from <https://phys.org/news/2019-05-d-technology-distant.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.