


New searches for supersymmetry presented by ATLAS experiment

May 23 2019



The ATLAS Experiment at CERN is searching for signs of supersymmetry, a theory which extends the Standard Model to answer many unsolved questions about the Universe. Credit: ATLAS Collaboration/CERN

The Standard Model is a remarkably successful but incomplete theory.

[Supersymmetry](#) (SUSY) offers an elegant solution to the Standard Model's limitations, extending it to give each particle a heavy "superpartner" with different [spin](#) properties (an important quantum number distinguishing matter particles from force particles and the Higgs boson). For example, sleptons are the spin 0 superpartners of spin 1/2 electrons, muons and tau leptons, while charginos and neutralinos are the spin 1/2 counterparts of the spin 0 Higgs bosons (SUSY postulates a total of five Higgs bosons) and spin 1 gauge bosons.

If these superpartners exist and are not too massive, they will be produced at CERN's Large Hadron Collider (LHC) and could be hiding in data collected by the [ATLAS detector](#). However, unlike most processes at the LHC, which are governed by strong force interactions, these superpartners would be created through the much weaker electroweak interaction, thus lowering their production rates. Further, most of these new SUSY particles are expected to be unstable. Physicists can only search for them by tracing their decay products—typically into a known Standard Model particle and the [lightest supersymmetric particle](#) (LSP), which could be stable and non-interacting, thus forming a natural dark matter candidate.

On 20 May, 2019, at the [Large Hadron Collider Physics](#) (LHCP) conference in Puebla, Mexico, and at the [SUSY2019](#) conference in Corpus Christi, U.S., the ATLAS Collaboration presented numerous new searches for SUSY based on the full LHC Run 2 dataset (taken between 2015 and 2018), including two particularly challenging searches for electroweak SUSY. Both searches target particles that are produced at extremely low rates at the LHC, and decay into Standard Model particles that are themselves difficult to reconstruct. The large amount of data successfully collected by ATLAS in Run 2 provides a unique opportunity to explore these scenarios with new analysis techniques.

Search for the "stau"

Collider and astroparticle physics experiments have set limits on the mass of various SUSY particles. However, one important superpartner—the tau slepton, known as the stau—has yet to be found beyond the exclusion limit of around 90 GeV found at the LHC's predecessor at CERN, the Large Electron-Positron collider (LEP). A light stau, if it exists, could play a role in neutralino co-annihilation, moderating the amount of dark matter in the visible universe, which otherwise would be too abundant to explain astrophysical measurements.




Figure 1: Left: Observed (expected) limits on the combined left and right stau pair production are shown by the red line (black dashed line). Right: Observed (expected) limits on the stau-left pair production are shown by the red line (black dashed line). The mass of stau is shown on the x-axis, while the mass of the LSP is shown on the y-axis. Credit: ATLAS Collaboration/CERN

The search for a light stau is experimentally challenging due to its extremely low production rate in LHC proton-proton collisions, requiring advanced techniques to reconstruct the Standard Model tau leptons it can decay into. In fact, during Run 1, only a narrow parameter

region around a stau mass of 109 GeV and a massless lightest neutralino could be excluded by LHC experiments.

This [first ATLAS Run 2 stau search](#) targets the direct production of a pair of staus, each decaying into one tau lepton and one invisible LSP. Each tau lepton further decays into hadrons and an invisible neutrino. Signal events would thus be characterised by the presence of two sets of close-by hadrons and large missing transverse energy (ET_{miss}) originating from the invisible LSP and neutrinos. Events are further categorised into regions with medium and high ET_{miss}, to examine different stau mass scenarios.

The ATLAS data did not reveal hints for stau pair production and thus new exclusion limits were set on the mass of staus. These limits are shown in Figures 1 using different assumptions on the presence of both possible stau types (left and right, referring to the two different spin states of the tau partner lepton). The limits obtained are the strongest obtained so far in these scenarios.

Compressed search

One of the reasons physicists have yet to see charginos and neutralinos may be because their masses are compressed. In other words, they are very close to the mass of the LSP. This is expected in scenarios where these particles are higgsinos, the superpartners of the Higgs bosons.




Figure 2: Observed (expected) limits on higgsino production are shown by the red line (blue dashed line). The mass of the produced higgsino is shown on the x-axis, while the mass difference to the LSP is shown on the y-axis. The grey region represents the models excluded by the LEP experiments; the blue region, the constraint from the previous ATLAS search for higgsinos. Credit: ATLAS Collaboration/CERN

Compressed higgsinos decay to pairs of electrons or muons with very low momenta. It is challenging to identify and reconstruct these particles in an environment with more than a billion high-energy collisions every second and a detector designed to measure high-energy [particles](#)—like trying to locate a whispering person in a very crowded and noisy room.

A [new search for higgsinos](#) utilises muons measured with unprecedentedly low—for ATLAS, so far—momenta. It also benefits from new and unique analysis techniques that allow physicists to look for higgsinos in areas that were previously inaccessible. For example, the search uses charged particle tracks, which can be reconstructed with very low momentum, as a proxy for one of the electrons or muons in the decay pair. Because of the small mass difference between the higgsinos, the mass of the electron/muon and track pair is also expected to be small.

Once again, no signs of higgsinos were found in this search. As shown in Figure 2, the results were used to extend constraints on higgsino masses set by ATLAS in 2017 and by the LEP experiments in 2004.

Overall, both sets of results place strong constraints on important supersymmetric scenarios, which will guide future ATLAS searches. Further, they provide examples of how advanced reconstruction techniques can help improve the sensitivity of new physics searches.

More information: Search for direct stau production in events with two hadronic tau leptons in 13 TeV proton-proton collisions with the ATLAS detector (ATLAS-CONF-2019-018):

[atlas.web.cern.ch/Atlas/GROUPS ... ATLAS-CONF-2019-018/](https://atlas.web.cern.ch/Atlas/GROUPS/CONF/PUBLISHED/2019/018/)

Searches for electroweak production of supersymmetric particles with compressed mass spectra in 13 TeV proton-proton collisions with the ATLAS detector (ATLAS-CONF-2019-014):

[atlas.web.cern.ch/Atlas/GROUPS ... ATLAS-CONF-2019-014/](https://atlas.web.cern.ch/Atlas/GROUPS/CONF/PUBLISHED/2019/014/)

Provided by ATLAS Experiment

Citation: New searches for supersymmetry presented by ATLAS experiment (2019, May 23)
retrieved 19 September 2024 from <https://phys.org/news/2019-05-supersymmetry-atlas.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.