


Microaerobic Fe(II) oxidation could drive microbial carbon assimilation in paddy soil

29 May 2019



Schematic diagram of microaerobic Fe(II) oxidation coupled to carbon assimilation processes driven by microbes from paddy soil. Credit: ©Science China Press

The carbon assimilation process driven by soil microorganisms is important to maintain the production and ecological function of paddy fields. A recent publication from Prof. Fangbai LI's group has found that Fe(II) oxidation under microaerobic conditions could drive microbial carbon assimilation in paddy soil, and identified the potential microaerophilic Fe(II)-oxidizing bacteria (FeOB). These findings provide an insight into the ecological function of iron cycling in the critical zone of red soil.

The research paper, "Microaerobic Fe(II) oxidation coupled to carbon assimilation processes driven by microbes from paddy soil," has been published in *Science China Earth Sciences* (2019, vol.62). The corresponding author of this study is Prof. Fangbai LI from Guangdong Institute of Eco-Environmental

Science & Technology, China. The first author is Prof. Xiaomin LI from SCNU Environmental Research Institute (SERI), South China Normal University (SCNU), China. The researchers employed gradient tubes and a ¹³C-labeled technique to investigate the processes of microaerobic Fe(II) oxidation coupled to carbon assimilation in the paddy soil from South China, and identified the associated microaerophilic FeOB.

Microbial growth using iron as substrate is believed to be an ancient form of metabolism, which likely evolved during the great oxidation event about 2.4 billion years ago, leading to banded iron formation (BIF). With a gradual increase in atmospheric O₂ concentrations over time, the habitats for microaerophilic FeOB became restricted to the iron-rich oxic-anoxic interfaces, such as groundwater, freshwater iron seeps, wetlands, sediments and marine hydrothermal vents. Most of the paddy soils in China are located in the southern region's red soils, where potential microaerobic conditions can be developed during the flooding season, such as the water-soil interface, and rice rhizosphere. However, few studies report such coupled processes in paddy soil of the critical zone in South China.

In this study, rhizosphere soil from flooded paddy fields was used as the inoculum to enrich the microaerophilic FeOB in gradient tubes with different Fe(II) substrates (FeS and FeCO₃) and ¹³C-biocarbonate as inorganic carbon source to track the carbon assimilation. Kinetics of Fe(II) oxidation and biomineralization were analyzed, and the composition and abundance of the microbial community were profiled using 16S rRNA gene-based high-throughput sequencing.

Results showed that microbial cell bands were formed 0.5-1.0 cm below the medium surface in the inoculated tubes with Fe(II) substances. The yields of ¹³C-biocarbonate incorporation was FeCO₃ (1.61-1.98 percent) > FeS (0.44-0.54 percent), and

the Fe(II) oxidation rate was FeS (0.156 mM d⁻¹) > FeCO₃ (0.106 mM d⁻¹). Relative to the agar only control, the abundances of Clostridium and Pseudogulbenkiania increased with FeS, while those of Vogesella, Magnetospirillum, Solitalea, and Oxalicibacterium increased with FeCO₃.

The results obtained in this study confirmed that microorganisms enriched from the paddy soil in the red soil area could couple microaerobic Fe(II) oxidation to carbon assimilation. Although it is difficult to evaluate the contribution of such coupling processes to iron mineral formation and carbon assimilation in the paddy field, such coupling processes driven by microorganisms should not be ignored, particularly during the iron-carbon transformation in the critical zone of red soils, given that several habitats potentially exist in paddy field for microaerophilic FeOB.

At the oxic-anoxic interface and rice rhizosphere, the microaerobic Fe(II) oxidation not only coupled [carbon](#) transformation, but also formed Fe(III) minerals with high surface area and reactivity, which can strongly influence the fates of heavy metals and other hazardous organic pollutants by adsorption, complexation, and redox reaction.

More information: Xiaomin Li et al, Microaerobic Fe(II) oxidation coupled to carbon assimilation processes driven by microbes from paddy soil, *Science China Earth Sciences* (2019). [DOI: 10.1007/s11430-018-9329-3](#)

Provided by Science China Press

APA citation: Microaerobic Fe(II) oxidation could drive microbial carbon assimilation in paddy soil (2019, May 29) retrieved 20 September 2021 from <https://phys.org/news/2019-05-microaerobic-feii-oxidation-microbial-carbon.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.