

Swapping water for CO₂ could make fracking greener and more effective

May 30 2019



A shale sample ready for fracturing with CO₂. Credit: Xuehang Song

Scientists at the Chinese Academy of Sciences and China University of Petroleum (Beijing) have demonstrated that CO₂ may make a better hydraulic fracturing (fracking) fluid than water. Their research, published May 30 in the journal *Joule*, could help pave the way for a more eco-friendly form of fracking that would double as a mechanism for storing captured atmospheric CO₂.

Fracking is a technique used to extract resources from unconventional reservoirs in which fluid (usually water mixed with sand, foaming agents, biocides, and other chemicals) is injected into the rock, fracturing it to release the resources within. Of the approximately 7-15 million liters of fluid injected, 30%-50% remains in the [rock formation](#) after extraction ends. Its high water consumption, [environmental risks](#), and frequent production issues have led to concerns about fracking among both industry experts and environmental advocates.

"Non-aqueous fracturing could be a potential solution to circumvent these issues," says Nannan Sun, a researcher in the Shanghai Advanced Research Institute at the Chinese Academy of Sciences. "We chose CO₂ fracturing from a range of options because the process includes multiple benefits. However, we were still lacking a fundamental understanding of the technology, which is greatly important for its further development and deployment."

Benefits of CO₂ fracturing include eliminating the need for a hefty water supply (which would make fracking viable in arid locations), reducing the risk of damage to reservoirs (as often happens when aqueous

solutions create blockages in the rock formation), and providing an underground repository for captured CO₂.

However, CO₂ is not likely to become commonly used as a fracking fluid unless it is more effective than water at resource production. To investigate the differences between CO₂ and water as fracturing fluids on a [microscopic level](#), Sun and his team collected shale outcrops from Chongqing, China and fractured them with both fluids. They found that CO₂ outperformed water, creating complex networks of [fractures](#) with significantly higher stimulated volumes.

"We demonstrated that CO₂ has higher mobility than [water](#), and, therefore, the injection pressure can be better delivered into the natural porosity of the formation," says Sun. "This changes the mechanism by which the fractures are created, generating more complex fracture networks that result in more efficient shale gas production."

While the researchers believe this hydraulic fracturing technology will be scalable, its large-scale development is currently limited by CO₂ availability. The cost of CO₂ captured from emission sources is still prohibitively expensive to make CO₂ an industry-wide [fracking](#) fluid replacement.

The team also notes that once CO₂ has been injected into the fracture, it acquires a low viscosity that inhibits it from effectively transporting sand to the fractures. Since the sand is intended to prop open the fractures while shale gas is harvested, it is critical that scientists learn to improve the fluid's viscosity—but the team is not yet sure how to do so while keeping costs low and minimizing the environmental footprint.

As next steps, the researchers plan to study the limits of CO₂ fracturing technology in order to better understand how it can be used. "Further investigations are needed to identify the effects of type of reservoirs,

geomechanical properties and conditions, CO₂ sensitivity of the formation, and so forth," says Sun. "Additionally, cooperation with industries will be carried out to push forward the practical deployment of the technology."

More information: *Joule*, Song, Guo, and Zhang et al.: "Fracturing with Carbon Dioxide: From Microscopic Mechanism to Reservoir Application" [www.cell.com/joule/fulltext/S2542-4351\(19\)30216-8](http://www.cell.com/joule/fulltext/S2542-4351(19)30216-8) , DOI: [10.1016/j.joule.2019.05.004](https://doi.org/10.1016/j.joule.2019.05.004)

Provided by Cell Press

Citation: Swapping water for CO₂ could make fracking greener and more effective (2019, May 30) retrieved 3 May 2024 from <https://phys.org/news/2019-05-swapping-co2-fracking-greener-effective.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.