

Scientists identify plant that flowers in Brazilian savanna one day after fire

June 18 2019, by José Tadeu Arantes



Rapid resprouting and flowering of *Bulbostylis paradoxa* is proof of the Cerrado biome's superb resilience and its capacity to evolve through fire. Credit: Alessandra Fidelis

Plants in the Brazilian savanna, the Cerrado, have evolved to deal with fire. When fire is used intelligently as part of a carefully planned land management method, it is indispensable to the conservation of this superb ecosystem, the world's most biodiverse savanna. Two months suffice for the Cerrado to burst into flower after a fire.

The study "From ashes to flowers: a savanna sedge initiates flowers 24 h

after [fire](#)", published recently in the journal *Ecology*, confirms this observation.

"The species is *Bulbostylis paradoxa*, a sedge [perennial herb similar to rushes and grasses] belonging to the family Cyperaceae," Alessandra Fidelis, first author of the article, told. "Its common name in Brazil is cabelo-de-índio ['Amerindian hair']."

Fidelis is a professor at São Paulo State University (UNESP) in Rio Claro, Brazil, and conducted the study with support from São Paulo Research Foundation—FAPESP as part of the project "How does the fire season affect Cerrado vegetation?".

The Cerrado is a unique type of savanna. Its capacity to grow back and flower after burning is a key difference in comparison to African and Australian savannas.

This had already been reported in the nineteenth and early twentieth centuries by naturalists who visited Brazil, such as French botanist Augustin de Saint-Hilaire (1779-1853) and Danish plant ecologist Eugen Warming (1841-1924).

Leopoldo Magno Coutinho (1934-2016), a professor at the University of São Paulo (USP), focused on it in his habilitation.

Fidelis has been studying postfire regeneration of the Cerrado since 2009, but in this study, she was struck by the speed with which *B. paradoxa* flowers after being completely charred in a wildfire. The article is indeed a first in this respect. "It's the only event of its kind described worldwide to date," she said.

B. paradoxa is distributed throughout South America and is found from Venezuela to the south of the continent. It flowers on a significant scale

only after fires. "In our experiments with burning as a careful management practice, we found that plants of this species reduced to carbonized stumps began exhibiting [white spots](#) 24 hours after the fire. These spots were incipient inflorescences," Fidelis recalled.

"In little more than a week, the [flowers](#) were fully formed and ready for pollination. The speed of this response is a significant advantage for the plant, enabling it to flower, fructify and disperse windborne seeds in the open, with the soil uncovered and without barriers or competitors. Only 40 days after a fire, it's very hard to find seeds because they've dispersed."

More generally, the abundant seeds available after fires in the Cerrado are an important resource for predators, such as ants and birds.

Resprouting leaves are tender and palatable to [large mammals](#), such as deer and cattle.

Burning is a problem when it results from arson or uncontrolled wildfires that spread disastrously owing to a buildup of combustible material after years without properly managed burning.

"The Cerrado has evolved through fire," Fidelis said. "Its vegetation regenerates easily. Species that did not occur before a fire may appear in certain areas. The animals may suffer losses, however, as many of them are trapped in fires. It is crucial to bear in mind that the Cerrado contains gallery forest, valley forest and palm marshes, where fire-sensitive species may fail to recover after a major burning. Hence, careful fire management is important. Preventive burning at the right time, with total area zoning and a rotating fire schedule for demarcated patches, is the best defense against disasters due to uncontrolled wildfire."

Expansion of the agricultural frontier, with large-scale monocultures and

the intensive use of machinery and herbicides, leaving the soil completely clear and subject to invasion by signalgrass and molasses grass, is currently the main threat to the Cerrado's survival. Improper use of fire is the next most important threat. Together, these two factors endanger the continued existence of the entire ecosystem.

Some of Brazil's most important rivers originate in the Cerrado, including the Xingu, Tocantins, Araguaia, São Francisco, Parnaíba, Gurupi, Jequitinhonha, Paraná and Paraguay rivers. In addition to the irreparable loss of biodiversity, destruction of the Cerrado jeopardizes the basins of these rivers as a major source of fresh water and hydropower potential.

More information: Alessandra Fidelis et al, From ashes to flowers: a savanna sedge initiates flowers 24 h after fire, *Ecology* (2019). [DOI: 10.1002/ecy.2648](https://doi.org/10.1002/ecy.2648)

Provided by FAPESP

Citation: Scientists identify plant that flowers in Brazilian savanna one day after fire (2019, June 18) retrieved 19 September 2024 from <https://phys.org/news/2019-06-scientists-brazilian-savanna-day.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.