


Black (nano)gold to combat climate change

July 3 2019



SHINE ON BLACK GOLD FOR A BRIGHTER FUTURE

When exposed to light, gold (Au) nanoparticles (NPs) generate catalysing "hot" electrons and "hotspots"

Au NP "Hot" electron

"Hotspot" Colloidosome

Black (nano)gold can be used for solar energy harvesting

To improve the catalyst's efficiency, "hotspots" in colloidosomes can be tuned by tuning the size and gaps between Au NPs

- CO₂ conversion
- Fuel CH₄
- Protein unfolding
- Aldehyde hydrosilylation
- Alcohol oxidation
- Seawater desalination

The use of black gold can help us move one step closer to combating climate change

Chemical Science PICK OF THE WEEK

Plasmonic colloidosomes of black gold for solar energy harvesting and hotspots directed catalysis for CO₂ to fuel conversion
Dhiman, Maity et al. (2019) | DOI: 10.1039/C9SC02369K

ROYAL SOCIETY OF CHEMISTRY

Use of black gold can get us one step closer to combat climate change. Credit: Royal Society of Chemistry, Chemical Science

Global warming is a serious threat to the planet and living beings. One of the main causes of global warming is the increase in the atmospheric CO₂ level. The main source of this CO₂ is from the burning of fossil fuels in our daily lives (electricity, vehicles, industry and many more).

Researchers at TIFR have developed the solution phase synthesis of

dendritic plasmonic colloidosomes (DPCs) with varying interparticle distances between the [gold](#) nanoparticles (NPs) using a cycle-by-cycle growth approach by optimizing the nucleation-growth step. These DPCs absorbed the entire visible and near-infrared region of solar light, due to interparticle plasmonic coupling as well as the heterogeneity in the Au NP sizes, which transformed gold material to black gold.

Black (nano)gold was able to catalyze CO₂ to methane ([fuel](#)) at atmospheric pressure and temperature, using solar energy. Researchers also observed the significant effect of the plasmonic hotspots on the performance of these DPCs for the purification of seawater to [drinkable water](#) via steam generation, temperature jump assisted protein unfolding, oxidation of cinnamyl alcohol using pure oxygen as the oxidant, and hydrosilylation of aldehydes.

The results were attributed to varying interparticle distances and particle sizes in these DPCs. Results indicate the synergistic effects of EM and thermal hotspots as well as hot electrons on DPCs performance. Thus, DPC catalysts can effectively be utilized as Vis-NIR light photocatalysts, and the design of new plasmonic nanocatalysts for a wide range of other chemical reactions may be possible using the concept of plasmonic coupling.

Raman thermometry and SERS (surface-enhanced Raman spectroscopy) provided information about the thermal and electromagnetic hotspots and local temperatures which was found to be dependent on the interparticle plasmonic coupling. The [spatial distribution](#) of the localized surface plasmon modes by STEM-EELS plasmon mapping confirmed the role of the interparticle distances in the SPR (Surface Plasmon Resonance) of the material.

Thus, in this work, by using the techniques of nanotechnology, the researchers transformed metallic gold to black gold, by changing the size

and gaps between [gold nanoparticles](#). Similar to trees, which use CO₂, sunlight and water to produce food, the developed black gold acts like an artificial tree that uses CO₂, sunlight and water to produce fuel, which can be used to run our cars. Notably, black gold can also be used to convert sea water into drinkable water using the heat that black gold generates after it captures sunlight.

This work is a way forward to develop 'artificial trees' to capture and convert CO₂ to fuel and useful chemicals. Although at this stage, the production rate of fuel is low, in coming years, these challenges can be resolved. We may be able to convert CO₂ to fuel using sunlight at atmospheric condition, at a commercially viable scale and CO₂ may then become our main source of clean energy.

More information: Mahak Dhiman et al, Plasmonic colloidosomes of black gold for solar energy harvesting and hotspots directed catalysis for CO₂ to fuel conversion, *Chemical Science* (2019). [DOI: 10.1039/C9SC02369K](#)

Provided by Tata Institute of Fundamental Research

Citation: Black (nano)gold to combat climate change (2019, July 3) retrieved 19 September 2024 from <https://phys.org/news/2019-07-black-nanogold-combat-climate.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.