


Ancient Saharan seaway shows how Earth's climate and creatures can undergo extreme change

July 8 2019



During the Late Cretaceous-early Paleogene, the shallow waters of the Trans-Saharan Seaway waters were teeming with aquatic species which ranged from small mollusks to giant sea snakes and catfish. Credit: © Carl Buell

A new paper to be published in the *Bulletin of the American Museum of Natural History* integrates 20 years of research by a diverse scientific team and describes the ancient Trans-Saharan Seaway of Africa that existed 50 to 100 million years ago in the region of the current Sahara Desert. Led by Maureen O'Leary, Professor of Anatomical Sciences at the Renaissance School of Medicine at Stony Brook University, the paper is a comprehensive synthesis and contains the first reconstructions of extinct aquatic species in their habitats along the seaway and places in context massive climate and sea level changes that can occur on Earth.

The region now holding the Sahara Desert was once under water, in striking contrast to the present-day arid environment. This dramatic difference in climate over time is recorded in the rock and fossil record of West Africa during a time range that extends through the Cretaceous-Paleogene (KPg) boundary. West Africa was bisected by a shallow saltwater body that poured onto continental crust during a time of high global sea level. The Bulletin paper involves an assessment and continued analysis of three expeditions led by Professor O'Leary (1999, 2003, and 2008) within rock exposures in the Sahara Desert in Mali, and subsequently the laboratory work of the fossil finds in the region.

"Fossils found on the expeditions indicate that the sea supported some of the largest sea snakes and catfish that ever lived, extinct fishes that were giants compared to their modern day relatives, mollusk-crushing fishes, tropical invertebrates, long-snouted crocodylians, early mammals and mangrove forests," explained Professor O'Leary, who is also a Research Associate in the Division of Paleontology, American Museum of Natural History. "Because the seaway changed in size and geography frequently, we propose that it may have resulted in 'islands of water' that stimulated species gigantism."

The paper contains the first reconstructions of ancient relatives of elephants and large apex predators such as sharks, crocodylians and sea

snakes.

"With our analysis and new technologies, such as a computer-aided map of the seaway, our work is an important step toward increasing our understanding of the KPg boundary event, the time of non-avian dinosaur extinction," said Professor O'Leary.

She and colleagues point out that the paper places in context climate and sea level changes that can occur on Earth.

For example, scientists currently predict that global warming will result in the sea rising two meters by the end of the 21st century. The study in the Bulletin describes how, in the Late Cretaceous, the time under study, sea level rise far exceeded that which is predicted by human-induced climate change. In the Late Cretaceous sea level was 300m higher than present—40 percent of current land was under water, which is very different from today. This information underscores the dynamic nature of Earth.

Professor O'Leary explained that scientists do not have detailed stratigraphic terrestrial/near shore sections with fossils on every continent to examine exactly how the KPg boundary unfolded globally. There is only one good nearshore or terrestrial section with vertebrate fossils in the western United States. The expeditions in Mali, she added, created a new section, which is imperfect, missing some of the earliest Paleogene yet contributes to a better understanding of global events 50 to 100 million years ago.

The expeditions spanning 20 years involved Professor O'Leary and numerous colleagues internationally to excavate the fossils and conduct the research. The collaborative research team consists of paleontologists and geologists from the United States, Australia and Mali.

"Few paleontologists had worked the region, given its remoteness and scorching 125 degree F temperatures. The shifting sand dunes made it difficult to find rocky outcrops, and worse still, a flash rain storm flooded the roadways making navigation nearly impossible," said Leif Tapanila, Ph.D., Professor of Geosciences at Idaho State University and a co-author of the paper. "These expeditions could not have succeeded without the experience of local Malian drivers and guides, and I was amazed by the quality and diversity of marine fossils we found in the Sahara Desert."

More information: Stratigraphy and paleobiology of the Upper Cretaceous-Lower Paleogene sediments from the Trans-Saharan Seaway in Mali. (Bulletin of the American Museum of Natural History, no. 436) digitallibrary.amnh.org/handle/2246/6950

Provided by American Museum of Natural History

Citation: Ancient Saharan seaway shows how Earth's climate and creatures can undergo extreme change (2019, July 8) retrieved 19 September 2024 from <https://phys.org/news/2019-07-ancient-saharan-seaway-earth-climate.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.