


On-demand control of terahertz and infrared waves

9 July 2019

Magneto-infrared microscope (made in Geneva)



The experimental device that focused infrared and terahertz radiation on small samples of pure graphene in the magnetic field, built by the UNIGE team. Credit: UNIGE, Ievgeniia Nedoliuk

The ability to control infrared and terahertz waves using magnetic or electric fields is one of the great challenges in physics that could revolutionise optoelectronics, telecommunications and medical diagnostics. A theory from 2006 predicts that it should be possible to use graphene—a monoatomic layer of carbon atoms—in a magnetic field not only to absorb terahertz and infrared light on demand but also to control the direction of circular polarisation. Researchers from the University of

Geneva (UNIGE), Switzerland, and the University of Manchester have succeeded in testing this theory and achieved the predicted results. The study, to be published in the journal *Nature Nanotechnology*, shows that the scientists found an efficient way to control infrared and terahertz waves. It also shows that graphene is keeping its initial promises, and is making its way as the material of the future, whether on earth or in space.

"There exist a class of the so-called Dirac materials, where the electrons behave as if they do not have a mass, similar to [light particles](#), the photons," explains Alexey Kuzmenko, a researcher at the Department of Quantum Matter Physics in UNIGE's Science Faculty, who conducted this research together with Ievgeniia Nedoliuk. One of such Dirac materials is [graphene](#), a monolayer of carbon atoms arranged in honeycomb structure, related to graphite used to make pencils.

The interaction between graphene and [light](#) suggests that this material could be used to control infrared and [terahertz waves](#). "That would be a huge step forward for optoelectronics, security, telecommunications and medical diagnostics," points out the Geneva-based researcher.

Backing up an old theory via experimentation

A theoretical prediction from 2006 posited that if a Dirac material is placed in a magnetic field, it will produce a very strong cyclotron resonance. "When a charged particle is in the magnetic field, it moves in a circular orbit and absorbs the [electromagnetic energy](#) at the orbiting, or cyclotron, frequency, as for example, it happens in the Large Hadron Collider at CERN," explains Alexey Kuzmenko. "And when the particles have charge but no mass, as electrons in graphene, the absorption of light is at its maximum!"

To demonstrate this maximum absorption, the physicists needed a very pure graphene so that the

electrons travelling long distances would not scatter on impurities or crystal defects. But this level of purity and lattice order are very difficult to obtain and are only achieved when graphene is encapsulated in another two-dimensional material—boron nitride.

The UNIGE researchers teamed up with the group from the University of Manchester led by André Geim—the 2010 Nobel Prize winner in Physics for discovering graphene—to develop extremely pure graphene samples. These samples, which were exceptionally large for this type of graphene, were nevertheless too small to quantify the cyclotron resonance with well-established techniques. This is why the Geneva researchers built a special experimental setup to concentrate the infrared and terahertz radiation on small samples of pure graphene in the [magnetic field](#). "And the result of the experiment confirmed the theory from 2006!" adds Alexey Kuzmenko.

apply magnetic fields, which could be generated already by inexpensive permanent magnets. Now that the theory has been confirmed, the researchers will continue to work on magnetically adjustable sources and detectors of [terahertz](#) and [infrared light](#). Graphene continues to surprise them.

More information: Ievgeniia O. Nedoliuk et al, Colossal infrared and terahertz magneto-optical activity in a two-dimensional Dirac material, *Nature Nanotechnology* (2019). [DOI: 10.1038/s41565-019-0489-8](#)

Provided by University of Geneva

Custom-controlled polarisation

The results demonstrated for the first time that a colossal magneto-optical effect occurs indeed if a layer of pure graphene is used. "The maximum possible magneto-absorption of the infrared light is now achieved in a monoatomic layer," says Kuzmenko.

In addition, the physicists found that it was possible to choose which circular polarisation—left or right—should be absorbed. "Natural or intrinsic graphene is electrically neutral and absorbs all light, regardless of its polarisation. But if we introduce electrically charged carriers, either positive or negative, we can choose which polarisation is absorbed, and this works both in the infrared and terahertz ranges," continues the scientist. This ability plays a crucial role, especially in the pharmacy, where certain key drug molecules interact with light depending on polarization direction. Interestingly, this control is considered promising for the search of life on exoplanets, since it is possible to observe the signatures of the molecular chirality inherent in biological matter.

Finally, the physicists found that to observe a strong effect in the terahertz range, it is sufficient to

APA citation: On-demand control of terahertz and infrared waves (2019, July 9) retrieved 27 February 2021 from <https://phys.org/news/2019-07-on-demand-terahertz-infrared.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.