

Teeth 'time capsule' reveals that 2 million years ago, early humans breastfed for up to 6 years

July 16 2019, by Renaud Joannes-Boyau, Ian Moffat, Justin W. Adams And Luca Fiorenza



The teeth in these *Australopithecus africanus* skulls contain important evidence about the nutrition of these individuals as they grew up. Credit: Luca Fiorenza, Author provided

Humans' distant ancestor *Australopithecus africanus* had a unique approach to raising their young, as shown in our new research published today in [*Nature*](#).

Geochemical analysis of four teeth shows they exclusively breastfed infants for about 6-9 months, before supplementing breast milk with varying amounts of solid food until they were 5-6 years old. The balance between milk and solid food in this period varied cyclically, probably in response to seasonal changes in food availability.

This knowledge is useful on several fronts. From an evolutionary point of view, it helps us understand the particular biological and behavioral adaptations of *Australopithecus africanus* compared to other extinct human ancestors and modern humans.

However, breastfeeding for up to 5-6 years is metabolically expensive—it requires a certain input of calories for the lactating mother. Using milk as a supplemental food for older offspring may have hampered the ability of the *A. africanus* species to successfully survive during a period of substantially changing climate.

Perhaps this way of life hastened the extinction of *A. africanus* around 2 million years ago.

A puzzling hominin

A. africanus was first discovered in 1924 by Australian-born scientist Raymond Dart at Taung in South Africa, and represented the first early human ancestor identified from Africa.



Illustration of a mother *Australopithecus africanus* and her young offspring.
Credit: Jose Garcia and Renaud Joannes-Boyau

A century of excavation and research later, Taung and other sites across South Africa produced a rich record of early human ancestors. This region is now a UNESCO World Heritage Site known as "[The Cradle of Humankind](#)".


This hominin species, a member of the human evolutionary lineage, had a mixture of ape-like characteristics and more specialized ones. It has only been recovered from fossil sites in South Africa that date to between 3 million and 2 million years ago.

Because only a few specimens exist, we have little information about how *A. africanus* lived and its relationship to other fossil hominin species such as the eastern African species of *Australopithecus*, the robust *Paranthropus*, and our own genus, *Homo*.

Zapping teeth

Our research takes advantage of cutting-edge analytical techniques. We used a laser to zap tiny pieces off fossil teeth, and then used an instrument called a mass spectrometer to determine their [chemical composition](#).

This is much less destructive than traditional methods that require the sample to be crushed and dissolved before analysis. This makes it a crucial technique for rare specimens such as those of *A. africanus*.



Schematic diagram of the use of laser ablation analysis to map the concentration of strontium and uranium within a tooth. Credit: Renaud Joannes-Boyau, Author provided

Our laser method also allowed us to map the composition of a specimen across the entire surface of a tooth—illuminating changes in diet, mobility or climate through time. This is an important advance, as it can reveal information that has been impossible to establish using conventional palaeontological methods.

In this study, we mapped changes in the concentration of barium, strontium and lithium in fossil teeth of two individuals. The amounts of these elements in our bodies can change significantly depending on our diet, and these changes are reflected in the composition of our bones and teeth.


While our bones continue to change composition as they remodel during

our lives, our teeth don't change after they form during childhood. Teeth are thus a perfect chemical time capsule of our childhood diet.

Mapping a varied diet

The concentration of barium in [breast milk](#) is very high, so infant teeth that form during breastfeeding will also have a high concentration of this element. This concentration [gradually drops as other sources of food are introduced](#).

The samples we analyzed from *A. africanus* show a different pattern, with cyclical fluctuations in barium concentration. This suggests mothers would increase or reduce the amount of additional food, probably depending on the availability of other resources. This is an adaptation to food stress also used by modern orangutans.



Australopithecus africanus canine showing a first period of nursing behaviour followed by a cyclical signal in the lithium, strontium and barium distribution.
Credit: Renaud Joannes-Boyau

The concentration of lithium in these teeth also varies cyclically, although not always at the same time as barium. The precise cause of lithium variations is still unclear but it seems to be linked to variations in body fat reserves or how much protein is eaten.


This suggests *A. africanus* regularly faced food stress, causing their diet and/or fat reserves to change with the seasons.

We compared the results from *A. africanus* to modern animals from similar savannah biome regions, which supported our results by showing cyclical signal linked to seasonal variations mix with another signal interpreted as cyclical breastfeeding also seen in modern orangutans.

Close to home

We also investigated the strontium isotope composition of these [teeth](#) to help understand where *A. africanus* was moving through the landscape. Isotopes of the same element can be distinguished by their mass.

Strontium isotopes are often used for this purpose in palaeontology, as different regions have characteristic isotope values that are taken up through food and drink.



Strontium isotopic ratio along the growth axis of an *Australopithecus africanus* tooth. Credit: Renaud Joannes-Boyau

The two *A. africanus* individuals in our study seemed to have lived most of their lives near the Sterkfontein cave where their remains were found.

Living in a region with limited food resources meant these early hominins would have eaten lots of different kinds of foods collected from varying habitats in order to survive.

Our research provides the first understanding of the nursing behavior of *A. africanus*. We now know this hominin had an extended period of breastfeeding supplemented by varying amounts of solid food that caused their fat reserves to fluctuate significantly.

This was likely part of a largely successful survival strategy for the species.

But as ecosystems changed with climate around 2 million years ago, the metabolic stress on mothers may have contributed to the eventual extinction of this species.

This article is republished from [The Conversation](#) under a Creative Commons license. Read the [original article](#).

Provided by The Conversation

Citation: Teeth 'time capsule' reveals that 2 million years ago, early humans breastfed for up to 6 years (2019, July 16) retrieved 19 September 2024 from <https://phys.org/news/2019-07-teeth-capsule-reveals-million-years.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.