

Global team of scientists finish assembling next-generation dark matter detector

August 7 2019, by Hayley Dunning



Researchers examine the foil-wrapped LUX-ZEPLIN xenon detector. Credit: Matt Kapust

The key component of the LUX-ZEPLIN experiment is ready to be sealed and lowered nearly 1.5 km underground, where it will search for

dark matter.


Dark matter is a mysterious form of matter thought to make up around 85 percent of the mass of the universe. However, because it is predicted to interact only very weakly with ordinary matter, it has so far not been detected.

LUX-ZEPLIN (LZ) will be the most sensitive dark matter experiment ever built. On 26 July, researchers finished assembling its centrepiece, the liquid [xenon](#) Time Projection Chamber (TPC), at the Sanford Underground Research Facility in South Dakota, U.S..

"This [xenon detector](#) will be at the heart of the LZ dark matter experiment," said Professor Henrique Araújo, from the Department of Physics at Imperial College London, who leads the LZ collaboration efforts in the UK and co-led the development of the TPC with Professor Tom Shutt from SLAC National Accelerator Laboratory.

13,500 hours of effort

To assemble the TPC, 250 members from 37 institutions from across the globe came together to ensure the mechanical, optical, electrical, radiological and cleanliness requirements of the project were met.



The recently assembled LUX-ZEPLIN xenon detector in the Surface Assembly Lab cleanroom at Sanford Underground Research Facility on 26 July 2019.
Credit: Matt Kapust.

Manufacture of the tens of thousands of components that make up the TPC started in 2015, and assembly of the instrument began in December 2018. The integration stage involved 13,500 hours of effort—a significant fraction of which was devoted to maintaining the required ultra-clean conditions at the surface-level assembly lab.

Next, it will be inserted into its cryostat vessel—a chamber that maintains cold temperatures—and lowered nearly 1.5km underground into a disused goldmine, ready to hopefully detect dark matter. Operations are planned to start in mid-2020.

"We have some things in common with a [space program](#)," said Professor Araújo. "Before you launch, you do all of your work on the ground for years, perfecting the engineering so your instrument will work no matter what. LZ is a bit like a space experiment, just headed the opposite direction. We cannot expose it to underground air—that would compromise its performance. Once you deploy it underground, that's it. It has to work."

Detecting WIMPs

Once underground, the detector will be cooled to -100°C and filled with ten tonnes of liquid xenon. Because xenon is a heavy element, there is a higher chance of xenon atoms interacting with hypothetical dark matter particles called WIMPs—weakly interacting massive particles.

Researchers believe that if a WIMP interacts with a xenon atom, it will produce two flashes of light. One appears promptly, when the particle collides with a xenon atom, which recoils through the liquid. The second is generated by electrons shaken off by the collision, which are guided to the top of the detector and accelerated through a layer of gaseous xenon above the liquid.



The detector is inspected under UV light. Credit: Nicolas Angelides

Although these flashes would be imperceptible to the human eye, the detector is lined with hundreds of photomultiplier tubes. These ultrasensitive sensors can amplify a signal from even a single photon of light.

Decades of development

The TPC design employed by LZ has been perfected over decades of experimentation with similar detectors that allow researchers to determine where an interaction happens, and whether it is likely due to a background interaction in the instrument or a true dark matter signal. Amongst these were the ZEPLIN-III experiment at the Boulby Mine in

North Yorkshire, led by Imperial, and the US-led LUX experiment—which gave rise to LZ.

Being buried deep underground protects the experiment from too many background interactions from processes readily detected at the surface that could obscure a dark [matter](#) signal, such as the cosmic rays that shower the Earth from outer space.

Professor Shutt said: "The TPC is a complex system and it's a major achievement to have it fully assembled. It takes us one important step closer to being able to look for [dark matter](#).

"It is also gratifying because it involved assembling a large number of subsystems designed and built by groups across the US and the UK over a number of years. So, it's a coming together of sorts for the collaboration."

Provided by Imperial College London

Citation: Global team of scientists finish assembling next-generation dark matter detector (2019, August 7) retrieved 20 September 2024 from <https://phys.org/news/2019-08-global-team-scientists-finish-next-generation.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--