

Northern tropical dry trend may just be normal variation

12 August 2019



Karst in southern Thailand Credit: Prof. Chuan-Chou Shen

Rainfall variations in the tropics not only potentially influence 40% of the world's population and the stability of the global ecosystem, but also the global hydrologic cycle and energy balance.

Beginning in the 20th century, a decline in northern tropical [rainfall](#) has been observed, with researchers unsure whether the decline stems from natural or anthropogenic causes.

New rainfall research has shed some light on this question, but left the final answer up in the air.

Recently, an international team led by Prof. Tan Liangcheng from the Institute of Earth Environment (IEE) of the Chinese Academy of Sciences developed a 2700-year rainfall record of the northern central Indo-Pacific region. The team's findings will be published in *PNAS*.

The researchers got official permission to collect stalagmites from the Klang Cave in southern Thailand. They used oxygen records and radiometric U-Th dating techniques with precision of half a year at best on three stalagmites to reveal regional rainfall history over the past 2700 years.

Results show millennial-scale decreasing regional rainfall, similar to other records from the northern tropics and opposite to southern tropical records.

This tropical interhemispheric precipitation seesaw pattern was most likely driven by changes in summertime insolation in each respective hemisphere. Notable centennial-decadal-scale dry (950-1150, 1200-1350, and 1910 to present) and wet (400-800) climate episodes occurred.

The extreme decadal wet event during the late 14th century and early 15th century coincided with extensive floods and destruction of the water management systems in Angkor, Cambodia, indicating the hydroclimatic impact on the decline of the Khmer Empire in the 15th century.

The scientists also developed a 2000-year Intertropical Convergence Zone (ITCZ) shift index record, which shows an overall southward shift over the past two millennia, and southward mean positions of ITCZ during the early Medieval Warm Period and the Current Warm Period in the central Indo-Pacific.

The scientists noted that the drying trend in the northern tropics since the early 20th century is similar to that from 950-1150 AD, which was attributed to a southward ITCZ shift and enhanced El Niño-like conditions. As a result, possible anthropogenic rainfall remains indistinguishable from natural variability in the northern tropics, Prof. Tan said. In other words, no clear case can be made that the recent northern tropical drying trend is caused by anthropogenic forces.

Although the cause of the last century's [drying trend](#) is uncertain, Prof. Tan sees a benefit in understanding historical rainfall trends.

"The rainfall variability during past warm periods can provide a historical analog to future trends under global warming, and help improve the

precision of climate models," Prof. Tan said.

More information: Liangcheng Tan et al.,
"Rainfall variations in central Indo-Pacific over the
past 2,700 y," *PNAS* (2019).

www.pnas.org/cgi/doi/10.1073/pnas.1903167116

Provided by Chinese Academy of Sciences

APA citation: Northern tropical dry trend may just be normal variation (2019, August 12) retrieved 13 April 2021 from <https://phys.org/news/2019-08-northern-tropical-trend-variation.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.