


Milky Way's black hole just flared, growing 75 times as bright for a few hours

August 14 2019, by Evan Gough



Four images from the paper. Over about a 2 hour period, Sgr A* flared to 75 times normal, and twice as bright as any other observed peak. At first, astronomers thought they were looking at the S star SO-2. Credit: Do et al; 2019.

Even though the black hole at the center of the Milky Way is a monster, it's still rather quiet. Called Sagittarius A*, it's about 4.6 million times more massive than the sun. Usually, it's a brooding behemoth. But scientists observing Sgr. A* with the Keck Telescope just observed its brightness blooming to over 75 times normal for a few hours.

The flaring is not visible in optical light. It's all happening in the near-infrared, the portion of the infrared spectrum closest to optical light. Astronomers have been watching Sgr. A* for 20 years, and though the black hole does have some variability in its output, this flaring event is like nothing astronomers have observed before. This peak was over twice as bright as the previous peak flux level.


These results are being reported in the *Astrophysical Journal Letters* in a paper titled "Unprecedented variability of Sgr A* in NIR," and is available at the prepress site arXiv.org. The lead author is Tuan Do, an astronomer at UCLA.

The team saw Sgr. A* flaring at 75 times normal for a two-hour period on May 13th. At first, [astronomer](#) Tuan Do thought that they were seeing a star called SO-2 rather than Sgr. A*. SO-2 is one of a group of [stars](#) called S-stars that orbits the black hole closely. Astronomers have been keeping an eye on it as it orbits the black hole.

Here's a timelapse of images over 2.5 hr from May from [@keckobservatory](#) of the supermassive black hole Sgr A*. The black hole is always variable, but this was the brightest we've seen in the infrared so far. It was probably even brighter before we started observing that night! pic.twitter.com/MwXioZ7twV

— Tuan Do (@quantumpenguin) [August 11, 2019](#)


In an interview with *ScienceAlert*, Do said, "The black hole was so bright I at first mistook it for the star S0-2, because I had never seen Sgr A* that bright. Over the next few frames, though, it was clear the source was variable and had to be the black hole. I knew almost right away there was probably something interesting going on with the black hole."



This is our best-yet image of an actual black hole. It's the super-massive black hole at the center of galaxy M87, and it was captured by the Event Horizon Telescope (EHT). The black hole itself can't actually be seen so this image is actually of its event horizon. The EHT's next target is Sgr. A*. Credit: Event Horizon Telescope Collaboration

The question is, what made Sgr. A* flare like this? At this point, astronomers aren't certain what caused the flaring. Sgr. A* has exhibited flaring before, just not as brightly. So flaring itself isn't unprecedented.

It's likely that something disrupted the black hole's usually quiet neighborhood, and there are at least a couple of possibilities. The first is not actually a disruption, but an inaccuracy in the statistical models used to understand the black hole. If that's the case, then the model needs to be updated to include these variations as "normal" for Sgr. A*.



The group of stars that orbit close to Sgr. A* are called S stars. SO-2 made it's closest approach about a year before the flaring observed in May 2019. Credit: Cmglee – Own work, CC BY-SA 3.0

The second possibility is where things get interesting: Something has changed in the black hole's neighborhood.


The previously mentioned star SO-2 is a prime candidate. It's one of two stars that approach very closely to Sgr. A* in an elliptical orbit. Every 16 years, it's at its closest. In the middle of 2018 was its last closest approach, when it was only 17 light-hours away from the black hole.

It's possible that SO-2's close approach disrupted the way that material flows into Sgr. A*. That would generate the kind of variability and bright flaring that astronomers saw in May, about one year after the star's close approach.

But astronomers aren't certain. SO-2 is not a very large star, and it seems unlikely that it could cause this type of disruption. Not only that, but it's the largest of the S stars that get close to Sgr. A*, so it's unlikely that one of the other stars could be the cause, either.

Another possibility is a [gas cloud](#). Back in 2002, astronomers saw what they thought might be a hydrogen gas cloud approaching the center of Sgr. A*. By 2012, astronomers were more certain that it was a cloud, and it was named G2. They measured the temperature of the cloud at 10,000 degrees Kelvin, and were able to measure its trajectory: In 2013, it would travel closely enough to the black hole that the tidal forces would tear it apart.

Initially, astronomers thought that gas from G2 might be drawn into Sgr. A*'s accretion disk, and that it would flare brightly as it was heated. But that never happened. But it's still possible that its passage close to the black hole set off a chain of events that caused or contributed to the May 2019 flaring.



A computer-simulated image of the hydrogen gas cloud G2 encountering Sgr. A* and being stretched out. The encounter could have disrupted the usually steady inflow of material into the black hole and caused the variability and flaring observed in May, 2019. Credit: M. Schartmann and L. Calcada/ European Southern Observatory and Max-Planck-Institut für Extraterrestrische Physik

In the final analysis (if there ever is one in science), this flaring may just be the natural result of a variable flow of material into Sgr. A*, which is expected to be lumpy. If that's the case, then we're back to updating the [statistical model](#) used to explain the black hole's variability.

The only way to know is to gather more data, not only with the Keck, while the galactic center is still visible at night, but with other telescopes. During the last few months, the [galactic center](#) has been visible, and 'scopes like the Spitzer, Chandra, Swift, and ALMA have been watching.

These observations across multiple wavelengths should help clarify the situation when they're made available.

More information: Unprecedented variability of Sgr A* in NIR, arXiv:1908.01777 [astro-ph.GA] arxiv.org/abs/1908.01777

Source Universe Today

Citation: Milky Way's black hole just flared, growing 75 times as bright for a few hours (2019, August 14) retrieved 25 April 2024 from <https://phys.org/news/2019-08-milky-black-hole-flared-bright.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.