

Total annihilation for supermassive stars

August 15 2019



Artist's concept of the SN 2016iet pair-instability supernova. Illustration by Joy Pollard. Credit: Gemini Observatory/NSF/AURA/

A renegade star exploding in a distant galaxy has forced astronomers to set aside decades of research and focus on a new breed of supernova that can utterly annihilate its parent star—leaving no remnant behind. The signature event, something astronomers had never witnessed before, may

represent the way in which the most massive stars in the Universe, including the first stars, die.

The European Space Agency's (ESA) Gaia satellite first noticed the supernova, known as SN 2016iet, on November 14, 2016. Three years of intensive follow-up observations with a variety of telescopes, including the Gemini North telescope and its Multi-Object Spectrograph on Maunakea in Hawai'i, provided crucial perspectives on the object's distance and composition.

"The Gemini data provided a deeper look at the supernova than any of our other observations," said Edo Berger of the Harvard-Smithsonian Center for Astrophysics and a member of the investigation's team. "This allowed us to study SN 2016iet more than 800 days after its discovery, when it had dimmed to one-hundredth of its peak brightness."

Chris Davis, program director at the National Science Foundation (NSF), one of Gemini's sponsoring agencies, added, "These remarkable Gemini observations demonstrate the importance of studying the ever-changing Universe. Searching the skies for sudden explosive events, quickly observing them and, just as importantly, being able to monitor them over days, weeks, months, and sometimes even years is critical to getting the whole picture. In just a few years, NSF's Large Synoptic Survey Telescope will uncover thousands of these events, and Gemini is well positioned to do the crucial follow-up work."

In this case, this deep look revealed only weak hydrogen emission at the location of the supernova, evidence that the progenitor star of SN 2016iet lived in an isolated region with very little star formation. This is an unusual environment for such a massive star. "Despite looking for decades at thousands of supernovae," Berger resumed, "this one looks different than anything we have ever seen before. We sometimes see supernovae that are unusual in one respect, but otherwise are normal; this

one is unique in every possible way."

SN 2016iet has a multitude of oddities, including its incredibly long duration, large energy, unusual chemical fingerprints, and environment poor in heavier elements—for which no obvious analogues exist in the astronomical literature.

"When we first realized how thoroughly unusual SN 2016iet is, my reaction was 'Whoa—did something go horribly wrong with our data?'" said Sebastian Gomez, also of the Center for Astrophysics and lead author of the investigation. The research is published in the August 15th issue of *The Astrophysical Journal*.




Image of SN 2016iet and its most likely host galaxy taken with the Low Dispersion Survey Spectrograph on the Magellan Clay 6.5-m telescope at Las Campanas Observatory in i-band on July 9, 2018. Credit: GEMINI Observatory

The unusual nature of SN 2016iet, as revealed by Gemini and other data, suggest that it began its life as a star with about 200 times the mass of our Sun—making it one of the most massive and powerful single star explosions ever observed. Growing evidence suggests the first [stars](#) born in the Universe may have been just as massive. Astronomers predicted that if such behemoths retain their mass throughout their brief life (a few million years), they will die as pair-instability supernovae, which gets its name from matter-antimatter pairs formed in the explosion.

Most massive stars end their lives in an explosive event that spews matter rich in [heavy metals](#) into space, while their core collapses into a neutron star or black hole. But pair-instability supernovae are a different breed. The collapsing core produces copious gamma-ray radiation, leading to a runaway production of particle and antiparticle pairs that eventually trigger a catastrophic thermonuclear explosion that annihilates the entire star, including the core.

Models of pair-instability supernovae predict they will occur in environments poor in metals (astronomer's term for elements heavier than hydrogen and helium), such as dwarf galaxies and the early Universe—and the team's investigation found just that. The event occurred at a distance of one billion light years in a previously uncatalogued dwarf galaxy poor in metals. "This is the first supernova in which the mass and metal content of the exploding star are in the range predicted by theoretical models," Gomez said.

Another surprising feature is SN 2016iet's stark location. Most [massive stars](#) are born in dense clusters of stars, but SN 2016iet formed in isolation some 54,000 [light years](#) away from the center of its dwarf host galaxy.

"How such a massive star can form in complete isolation is still a mystery," said Gomez. "In our local cosmic neighborhood, we only know

of a few stars that approach the mass of the star that exploded in SN 2016iet, but all of those live in massive clusters with thousands of other stars." To explain the event's long duration and slow brightness evolution, the team advances the idea that the progenitor star ejected matter into its surrounding environment at a rate of about three times the mass of the Sun per year for a decade before the star blew itself into oblivion. When the star ultimately exploded, the supernova debris collided with this material powering SN 2016iet's emission.

"Most supernovae fade away and become invisible against the glare of their host galaxies within a few months. But because SN 2016iet is so bright and so isolated we can study its evolution for years to come," said Gomez. "The idea of pair-instability supernovae has been around for decades," said Berger. "But finally having the first observational example that puts a dying star in the right regime of mass, with the right behavior, and in a metal-poor dwarf galaxy is an incredible step forward."

Not long ago, it was not known if such supermassive stars could actually exist. The discovery and follow-up observations of SN 2016iet have provided clear evidence for their existence and potential for affecting the development of the early Universe. "Gemini's role in this amazing discovery is significant," said Gomez, "as it helps us to better understand how the early Universe developed after its 'dark ages'—when no star formation occurred—to form the splendor of the Universe we see today."

More information: Sebastian Gomez et al. SN 2016iet: The Pulsational or Pair Instability Explosion of a Low-metallicity Massive CO Core Embedded in a Dense Hydrogen-poor Circumstellar Medium, *The Astrophysical Journal* (2019). [DOI: 10.3847/1538-4357/ab2f92](https://doi.org/10.3847/1538-4357/ab2f92) , arxiv.org/abs/1904.07259

Provided by Gemini Observatory

Citation: Total annihilation for supermassive stars (2019, August 15) retrieved 20 September 2024 from <https://phys.org/news/2019-08-total-annihilation-supermassive-stars.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.