


Genetic diversity couldn't save Darwin's finches

August 21 2019, by Michael Miller



A three-dimensional computer animation helps researchers compare the morphology of different finch species. Credit: Jose Barreiro

A study by the University of Cincinnati found that Charles Darwin's famous finches defy what has long been considered a key to evolutionary success: genetic diversity.

The research on finches of the Galapagos Islands could change the way

conservation biologists think about a species' potential for extinction in naturally fragmented populations.

UC graduate Heather Farrington and UC biologists Kenneth Petren and Lucinda Lawson found that genetic diversity was not a good predictor of whether populations of finches would survive. The study was published in *Conservation Genetics*.

A UC lab analysis of century-old museum specimens found that six of eight extinct populations had more genetic diversity than similar museum specimens from which descendants survive today. In most other species, low genetic diversity is a signal of a [population](#) in decline.

Researchers examined 212 tissue samples from museum specimens and living birds. Some of the museum specimens in the study were collected by Darwin himself in 1835. Only one of the extinct populations, a species called the vegetarian finch, had lower genetic diversity compared to modern survivors.

Lawson said the findings are explained by the fact that these birds can migrate in between populations.

Specifically, researchers believe a biological phenomenon called sink-source dynamics is at play in which larger populations of birds from other islands act as a "source" of immigrants to the island population that is naturally shrinking, the "sink." Without these immigrant individuals, the natural population on the island likely would continue to dwindle to local extinction. The immigrants have diverse genetics because they are coming from a variety of healthier islands, giving this struggling "sink" population inflated genetic diversity.

Petren said the findings serve as a warning that the genetics of individuals in fragmented populations might not tell the whole story

about a species. And that is important for scientists who increasingly use genetics to account for the flow of genes between populations when determining a threatened species' likelihood of extinction.

"The promise of genetics is to sample a few individuals to understand the whole population. But it's a cautionary note that you might be sampling a fragment. You could be misled," he said.



A small tree finch. Credit: Jack Stenger

Petren has been studying the birds for 25 years at UC's McMicken College of Arts and Sciences. He said the island's 18 recognized species of finches are unusual for other reasons. Some finches that look most different are actually closely related, he said. And similar-looking finches that birders might have trouble telling apart are actually far apart on the evolutionary family tree.

"It's a paradox. If Darwin fully understood what was going on, it might have blown his mind," Petren said. "These finches are not the first case you would pick to formulate the notion that species can change over time because the patterns of change are so complicated."

The UC study suggests that genetic diversity may not be the best predictor of extinction risk for mobile species like the island-hopping finches. That's because healthier populations may contribute individuals to declining ones.

Lawson said factors such as historical diversity or the possibility of gene flow between populations should be considered in addition to the snapshot view provided by a genetic analysis for a fuller understanding of a species' potential for extinction.

"Typically, we would expect populations with high genetic diversity to have a greater potential for long-term survival," she said. "Meanwhile, the low-diversity populations would be more likely to go extinct because that's a common pattern as populations decline to few individuals. Surprisingly, we found that most of the extinct populations had higher [genetic diversity](#)."

Extinction is a natural process in the crucible of life, but the impacts of people are increasingly apparent. Conservation biologists are developing [new tools](#) to identify species at risk of extinction before it is too late to save them. Scientists in the Galapagos are monitoring the finches to ensure their future survival. But researchers still don't know what drives some populations to extinction.

Lawson said many factors contribute to local or total extinctions. An island's size can influence local extinction potential. Small islands offer fewer resources and habitat. Likewise, how generalized or specialized an animal is plays a part as well. Species that rely on very specific habitats or food requirements are more vulnerable to extinction than generalists, for example. And habitat disturbances from human impacts, competitors or invasive species also affect survival.

The study was sponsored in part by the National Science Foundation, Sigma Xi, the American Ornithologists' Union and UC's Office of Research.

In a second related UC study, published in the journal *Ecography*, Lawson, Petren and former UC postdoctoral fellow John Niedzwiecki sought to better quantify movement for the finches within the archipelago to better understand the dynamics of gene flow which so significantly impact extinction potential.



UC doctoral graduate Jack Stenger conducts fieldwork on finches on the Galapagos island of San Cristobal. Credit: University of Cincinnati

Using mist nets, researchers captured 1,190 birds representing 11 species and 13 of the archipelago's islands. A tiny blood sample was taken from each bird before their release.

Scientists have several models explaining how animals spread to new islands. Generally, large islands provide immigrants to smaller islands. Likewise, older islands supply immigrants to newer ones over time.

UC researchers looked for a pattern of movement to test these theories in the Galapagos finches. They found that the finches flew against prevailing winds from the smallest outer islands to the biggest central ones, counter to hypotheses. The new arrivals might have been drawn to bigger islands they could see from afar, Lawson said.

"There is a visual cue. Those central islands are so much higher and capture the rain clouds," she said. "So if you take off, you might see those central islands and say, 'Land!'"

Galapagos finches are mediocre flyers. They don't have the physical stamina to leave the archipelago. And what's worse for birds surrounded by a vast ocean, their feathers are not particularly waterproof.

"If they get hit by a rogue wave, they sink. They're not well adapted to long-distance flights," Lawson said.

Conditions on the smallest islands can be especially hostile, with little water and fewer sources of food and cover. In his field notes, Darwin described one such island as "black, dismal heaps of broken lava, forming a shore fit for Pandemonium." Birds that evolved to cope with these extreme conditions would find comparatively greener pastures on the bigger central islands.

"On the smaller islands, you'd have to be a tough bird to make it. You really need to be tightly adapted to the resources to survive," Lawson said. "If you're a soft central archipelago bird and you try to make it on an island like Pinta, you'll probably die."

Darwin's *On the Origin of Species* was groundbreaking in our understanding of evolution through natural selection. "Survival of the fittest" is a household phrase and a shorthand description of any competition.



The feather of a Galapagos finch. Credit: Joseph Fuqua II/UC Creative Services

While scientists today know more about how new species are formed, the principles Darwin developed remain the foundation of evolutionary

biology, Petren said.

"Certainly, genetics is new. But the fundamental principles still stand. It's amazing how much of his work remains fundamentally true today," Petren said.

While scientists now understand a lot about evolution in the Galapagos Islands, and Darwin's finches in particular, there is still more to learn, Lawson said.

Petren and Lawson are working with UC graduate student Jose Barreiro, a native of Ecuador, to use advanced analytic tools to capture 3-D shapes of the finches' beaks to understand the selective pressures on shape that have led to this rapid radiation of species. This technique allows for a more precise comparison of an animal's features than previously was possible with callipers and other mechanical means.

Barreiro first visited the Galapagos on a class trip as a child.

"I think the Galapagos occupies a very special place in Ecuadorian identity," he said.

UC's researchers said the World Heritage Site is special to them as well. Petren led a multidisciplinary Honors class to the Galápagos to study sustainability on the islands. Now Petren and Lawson hope to create a similar experience to examine the islands from scientific, cultural and artistic perspectives.

"What's cool about the Galapagos is it sparks something in people. Of course, it's interesting for its geology and biology," Lawson said.

"But the islands have a lot more to offer than scientific understanding. The Galapagos have their own sounds, their own colors, their own

shapes—which could be inspiring for music, art, dance and any creative interpretations of these small desert islands in the middle of the ocean. It would be fantastic to get a diverse group of people to tour and visit the [islands](#) and represent UC's strengths to come up with something collaborative."

More information: Heather L. Farrington et al. Predicting population extinctions in Darwin's finches, *Conservation Genetics* (2019). [DOI: 10.1007/s10592-019-01175-3](#)

Lucinda P. Lawson et al. Darwin's finches: a model of landscape effects on metacommunity dynamics in the Galápagos Archipelago, *Ecography* (2019). [DOI: 10.1111/ecog.04511](#)

Provided by University of Cincinnati

Citation: Genetic diversity couldn't save Darwin's finches (2019, August 21) retrieved 19 September 2024 from <https://phys.org/news/2019-08-genetic-diversity-couldnt-darwin-finches.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--