

Global warming may diminish plant genetic variety in Central Europe

August 29 2019



The researchers have investigated the adaptability of *Arabidopsis* plants to climate change in greenhouses. The species will lose its genetic diversity under the increasingly dry climate around the Mediterranean. Credit: MPI f. Developmental Biology/ M. Exposito-Alonso

Plant genetic varieties in Central Europe could collapse due to temperature extremes and drought brought on by climate change. According to a new paper, only a few individuals of a species have

already adapted to extreme climate conditions. These findings suggest that the overall species genetic diversity could be greatly diminished. The publication was led by Moises Exposito-Alonso, who joins Carnegie next month from the Max Planck Institute for Developmental Biology and the University of California Berkeley.

An international team of researchers from the Max Planck Institute for Developmental Biology, University of Tübingen, Technical University of Madrid, and UC Berkeley studied populations of the thale cress plant, *Arabidopsis thaliana*, collected from over 500 geographic locations in Europe, commonly used for biological research. Growing these plants in Spain and Germany under dry conditions revealed how individual plants responded to heat and drought.

The investigators were particularly interested in how the unique blend of genetic mutations enables the [different individuals](#) of the same species to resist experimentally simulated climates. As some of these mutations may confer physiological advantages, the main goal of this study was to rank their fitness for the future survival of the species.

Models forecast shrink potential of European biodiversity

This data was then combined with models predicting how temperatures and precipitation are expected to shift geographically in the next few decades in order to understand how plant biodiversity will be affected by [climate change](#) caused by human activity.

"On the basis of our calculations up to the year 2050, we can determine a significant change in the mutations that will be needed for the thale cress to survive in Southern to Central Europe," first author Moises Exposito-Alonso said. "It is remarkable how much individuals from different parts

of Europe differ in their ability to withstand future climate conditions," added Detlef Weigel, Director at the Max Planck Institute, where the work was coordinated.

Many plant populations won't survive

As precipitation decreases and temperatures rise, especially in so-called transition zones between the Mediterranean and northern Europe, the team's predictions indicate that many of the continent's predominant plant populations will not possess the necessary genetic mutations to survive. These patterns might be shared across many plant species of Europe. While [genetic information](#) for most species is still lacking, the rapid advance in modern genetic methods allows researchers to obtain such information for more and more species. With such information in hand, it will be possible to improve predictions of where a species is most at risk of suffering from the consequences of climate change.

More information: Natural selection on the *Arabidopsis thaliana* genome in present and future climates, *Nature* (2019). [DOI: 10.1038/s41586-019-1520-9](#)

Provided by Max Planck Society

Citation: Global warming may diminish plant genetic variety in Central Europe (2019, August 29) retrieved 22 September 2024 from <https://phys.org/news/2019-08-global-diminish-genetic-variety-central.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
