

Mathematical model could help correct bias in measuring bacterial communities

September 10 2019, by Tracey Peake



Credit: NC State

Researchers from North Carolina State University have developed a mathematical model that shows how bias distorts results when measuring bacterial communities through metagenomic sequencing. The proof-of-concept model could be the first step toward developing calibration methods that could make metagenomic measurements more accurate.

Metagenomic sequencing identifies the number and type of [bacteria](#) present in a particular community—for example, in a human gut microbiome—through DNA extracted from the sample. "We're measuring communities of bacteria—which ones are present and how many of each one are there," says Ben Callahan, assistant professor of population health and pathobiology and corresponding author of a paper describing the work. "However, the measurement technology isn't perfect, which introduces [bias](#) into the results. And that means we don't get an accurate picture of the community we're trying to measure."

According to Callahan, since metagenomic sequencing is a multi-step process, biases can be introduced in every step.

"The most well-known step is DNA extraction, where we break open the bacteria to get to the DNA," Callahan says. "The cells of some bacteria are harder to break open than others. Let's say I have a bacterium that makes up half of the community but doesn't break very well. I could end up with only 10% of this bacterium in my measurement, instead of the 50% that is actually there. That introduces bias. Now every measurement or calculation I do from that point onward is systematically skewed."

Callahan, with NC State postdoctoral researcher Michael McLaren and biostatistician Amy Willis from the University of Washington, tested their [model](#) of bias against two types of [metagenomic sequencing](#)—16S RNA gene and shotgun metagenomics—in microbial communities of known composition, and found that the model accurately described bias in those circumstances.

"What this experiment shows is that the model we propose works in at least these limited circumstances," Callahan says. "The long-term goal is to provide a calibration tool for metagenomic measurements of complex natural communities, just as we have standards that we use to calibrate measurement technologies like scales, oscilloscopes and microscopes."

This work is a first step toward that."

The research appears in *eLife*. McLaren is first author.

More information: Michael R McLaren et al, Consistent and correctable bias in metagenomic sequencing experiments, *eLife* (2019).
[DOI: 10.7554/eLife.46923](https://doi.org/10.7554/eLife.46923)

Provided by North Carolina State University

Citation: Mathematical model could help correct bias in measuring bacterial communities (2019, September 10) retrieved 19 September 2024 from <https://phys.org/news/2019-09-mathematical-bias-bacterial.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.