

Scientists find ways to improve cassava, a 'crop of inequality' featured at Goalkeepers

September 25 2019



Today, University of Illinois scientist Amanda De Souza highlighted a crop of inequality called cassava at the Bill & Melinda Gates Foundation's third annual Goalkeepers event in New York City. Recently, De Souza and colleagues published a study in *New Phytologist* that identified opportunities to improve cassava yields -- which have not increased for more than fifty years. Credit: Claire Benjamin/RIPE Project

Today, as world leaders gather for the UN General Assembly, hundreds of emerging leaders focused on fighting global inequality came together at the Bill & Melinda Gates Foundation's third annual [Goalkeepers](#) event in New York City. Among them, University of Illinois scientist [Amanda De Souza](#) highlighted a crop of inequality called cassava, which has starchy, tuberous roots that sustain more than 500 million people in sub-Saharan Africa, yet cassava has been largely neglected by research and development compared to the staple crops of wealthier regions. Recently, De Souza and a team from Realizing Increased Photosynthetic Efficiency (RIPE) published a study in *New Phytologist* that identified opportunities to improve cassava yields—which have not increased for more than fifty years in Africa.

"For smallholder farmers who depend on tiny plots of land to feed and support their families, cassava is a 'backup' crop when other [crops](#) fail," De Souza said at Goalkeepers, where she described her work to improve cassava through the RIPE project. "Especially for women, who represent a majority of [smallholder farmers](#), cassava is a savings account. It is a resource they can harvest all year to pay for things like medical treatments and their children's school fees."

The RIPE project is an [international effort](#) to develop more productive crops by improving photosynthesis—the natural, sunlight-powered process that all plants use to fix [carbon dioxide](#) into carbohydrates that fuel growth, development, and ultimately yields. RIPE is supported by the Gates Foundation, the U.S. Foundation for Food and Agriculture Research (FFAR), and the U.K. Government's Department for International Development (DFID).

Led by RIPE researchers at Illinois and Lancaster University, this study examined factors that limit photosynthesis in 11 popular, or farmer-preferred, African varieties of cassava with the goal to eventually help cassava overcome photosynthetic limitations to boost yields.

First, the team examined the photosynthetic limitations of cassava exposed to constant high levels of light, like a plant would experience at midday with cloudless skies. In these conditions, and like many crops, cassava's photosynthesis is limited (by as much as 80 percent) by two factors: One half is due to the low speed that carbon dioxide molecules travel through the leaf to reach the enzyme that drives photosynthesis, called Rubisco. The other half is because Rubisco sometimes fixes oxygen molecules by mistake, wasting large amounts of the plant's energy.

Next, the team evaluated the limitations of photosynthesis under fluctuating [light conditions](#). Surprisingly, and unlike most crops, Rubisco was not the primary limiting factor when leaves transitioned from shade to sunlight, like when the sun comes out from behind a cloud. Instead, cassava is limited by stomata, which are microscopic pores on the surface of leaves that open to allow carbon dioxide to enter the plant but at the cost of water that escapes through these same pores. Stomata are partially closed in the shade and open in response to light when Rubisco is active.

"Rubisco is the major limiting factor during this transition from shade to light for most plants, including rice, wheat, and soybean," De Souza said. "Cassava is the first crop that we have found where stomata limit photosynthesis during these light transitions more than Rubisco."

Illinois' Postdoctoral Researcher Yu Wang created a computer model to quantify how much cassava would gain by overcoming this limitation. According to the leaf-level model, if stomata could open three times faster, cassava could fix 6 percent more carbon dioxide each day. In addition, cassava's water use efficiency—the ratio of biomass produced to water lost by the plant—could be improved by 16 percent.

In addition, the team found that it takes as long as 20 minutes for cassava

to transition from shade to full light and reach the maximum rate of photosynthesis, which is quite slow compared to other crops such as rice that can transition in just a few minutes. However, the fastest variety of cassava could transition almost three times faster and fix 65 percent more carbon dioxide into carbohydrates than the slowest variety. Closing this gap is another opportunity to improve cassava's productivity.

"Plants are constantly moving from shade to light as leaves shift and clouds pass overhead," said RIPE Director [Stephen Long](#), Ikenberry Endowed University Chair of Crop Sciences and Plant Biology at Illinois' Carl R. Woese Institute for Genomic Biology, who contributed to this study. "We hope that the variation that we discovered during these light transitions among cassava varieties can be used to identify new traits, and therefore opportunities for us to improve [cassava's](#) photosynthetic efficiency and yield potential."

More information: Amanda P. De Souza et al, Photosynthesis across African cassava germplasm is limited by Rubisco and mesophyll conductance at steady-state, but by stomatal conductance in fluctuating light, *New Phytologist* (2019). [DOI: 10.1111/nph.16142](https://doi.org/10.1111/nph.16142)

Provided by University of Illinois at Urbana-Champaign

Citation: Scientists find ways to improve cassava, a 'crop of inequality' featured at Goalkeepers (2019, September 25) retrieved 27 April 2024 from <https://phys.org/news/2019-09-scientists-ways-cassava-crop-inequality.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--