


# Researchers find antibiotic resistant genes prevalent in groundwater

October 4 2019



A representation of the study's sampling points (blue text) across (1) an advanced water treatment facility (AWTF) train and (2) the groundwater aquifer which receives its product water through percolation basins. Credit: Adam Smith, USC Viterbi School of Engineering

With climate change comes increasing water shortages, and potentially

longer periods of drought. As policymakers look urgently to wastewater recycling to stem the gap in water resources, the question is— how best to reuse water and ensure public safety. New and emerging contaminants like antibiotic resistant genes (ARGs) pose a potential hazard to public safety and water security. One concern is the spread of ARGs through the water system and an increase in development of antibiotic-resistant super bugs.


Adam Smith, Assistant Professor in the Sonny Astani Department of Civil and Environmental Engineering at the University of Southern California, and a team of researchers including Moustapha Harb, an Assistant Professor at Lebanese American University and USC Viterbi School of Engineering Ph.D. students Phillip Wang and Ali Zarei-Baygi, studied and compared samples from an advanced groundwater treatment facility in Southern California and groundwater aquifers to detect differences in ARG concentrations. While they found that the advanced groundwater treatment facility reduced nearly all targeted ARGs to below detection limits, groundwater samples had a ubiquitous presence of ARGs in both control locations and locations recharged with water from the advanced water treatment facility.

Historically, indirect reuse treatment methods in which an environmental barrier is an intermediary step in the water cleaning process have been more popular than the direct "toilet to tap" process. While indirect methods of water reuse treatment were, from a public perception and appetite, considered more reliable, it is actually direct reuse "toilet to tap" approaches which do not introduce an environmental buffer that produce safer, more pure water for potability. The reason for this lies in the way ARGs in the environment can contaminate potable reuse water. These findings were highlighted in a study published in *Environmental Science & Technology Letters*.

## **How ARGs Spread through Water Treatment**

## Systems

While some ARGs are naturally occurring in microbial communities, antibiotics, ARGs and antibiotic resistant pathogens are on the rise in water sources as a result of the overuse of antibiotics in general. In a typical water treatment cycle, wastewater is treated first at a wastewater treatment facility. The study found that this water remains high in ARGs, as they persist throughout the treatment process. From here, water intended for potable reuse is further purified using advanced physical and chemical techniques including reverse osmosis—a process that uses a partially permeable membrane to purify drinking water.



This chart shows ARG levels (normalized per liter of water sample) for (A) the advanced water treatment facility (AWTF) train ("P") samples and (B) groundwater ("G") samples. Credit: Adam Smith, USC Viterbi School of Engineering

In an indirect reuse schema, the purified water will be infused back into an environmental buffer, like a groundwater aquifer. Later, water is pulled from the aquifer and further treated at a drinking water treatment plant before being added to the public water supply. In contrast, in direct reuse approaches, purified water does not return to an environmental buffer, but instead, remains within the engineered water cycle, going from the [wastewater treatment plant](#) to the water reuse plant to the drinking water treatment plant and then out to your tap

Looking at the differences in ARGs between various water sources is incredibly important in considering future health hazards, like development of super bugs, said Smith. Since wastewater treatment plants are not generally designed for removal of micropollutants like [antibiotics](#), they tend to persist in treatment systems, leading to high densities of ARG resistant bacteria at different stages of treatment. When this water is introduced into an aquifer, where ARGs are already naturally occurring, it can become contaminated with ARGs and antibiotic resistant bacteria. To further complicate the issue, ARGs are easily transferred through horizontal gene transfer, increasing the risk for antibiotic resistant pathogens.

"ARGs are not regulated in any way and are a challenging emerging contaminant of concern due to our reliance on biological treatment in the engineered water cycle," Smith said. "Because they are biological contaminants—small fragments of DNA that are released to the environment—bacteria present in receiving environments can uptake them, becoming resistant themselves, and further perpetuating the spread of resistance."

Wastewater reuse is the prevailing option for dealing with a mounting pressure on global water supply and might be preferable to options like desalination, which is expensive and energy inefficient by comparison. However, the danger of spreading antibiotic resistance is one that should

inform which methodologies gain more traction and investment as we look ahead Smith said. Eliminating unknowns that persist in the environmental water buffers could be one way to ensure [water](#) that reaches our taps is clean of ARGs and other harmful contaminants.

"Lessening the global spread of antibiotic resistance will require an interdisciplinary approach that spans environmental and clinical systems. We must act fast before we enter a so called 'post-antibiotic world' where bacterial infections become impossible to treat," Smith said.

**More information:** Moustapha Harb et al, Background Antibiotic Resistance and Microbial Communities Dominate Effects of Advanced Purified Water Recharge to an Urban Aquifer, *Environmental Science & Technology Letters* (2019). [DOI: 10.1021/acs.estlett.9b00521](https://doi.org/10.1021/acs.estlett.9b00521)

Provided by University of Southern California

Citation: Researchers find antibiotic resistant genes prevalent in groundwater (2019, October 4) retrieved 23 April 2024 from <https://phys.org/news/2019-10-antibiotic-resistant-genes-prevalent-groundwater.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.