


Ultrafast particle interactions could make quantum information devices feasible

October 17 2019, by José Tadeu Arantes



Knowledge could contribute to the development of materials that prolong the coherence time. Credit: FAPESP

Energy is information. Lengthening the time during which a system is capable of retaining energy before losing it to the local environment is a key goal for the development of quantum information. This interval is called the "coherence time." Several studies have been performed with the aim of retarding decoherence.

A study conducted by researchers at the University of Campinas's Gleb Wataghin Institute of Physics (IFGW-UNICAMP) in São Paulo State, Brazil, and international collaborators set out to understand the decoherence process on the femtosecond (10^{-15} s) timescale. An article

describing the results was published in *Physical Review Letters*.

In the study, interactions between excitons (excited electrons) and phonons (quantum units of vibrational [energy](#) in a crystal lattice) were observed on the femtosecond timescale. A femtosecond is one quadrillionth of a second.

The use of a revolutionary ultrafast spectroscopy technique with high temporal and [spectral resolution](#) was fundamental to the success of the study. Lázaro Aurélio Padilha Jr. was one of the [principal investigators](#) for the project, and Diogo Burigo Almeida, then a postdoctoral fellow at Michigan, was one of the main authors. The experiment was performed with semiconducting nanocrystals dispersed in a colloidal solution at cryogenic temperatures.

"We found that when the material is excited [by light], the light it emits changes color in under 200 femtoseconds. This is due to interaction between excitons and phonons. The excitons transfer part of the energy they receive to the crystal lattice. This causes a change of frequency and hence a change of emission color," Padilha said.

Their study was the first to observe this phenomenon. "It had never been observed before because the amount of energy transferred from each exciton to the lattice is tiny, corresponding to 26 millielectron volts (26×10^{-3} eV), and the process takes a very short time, lasting under 200 femtoseconds (200×10^{-15} s). Similar phenomena have been observed but on far greater timescales and due to other processes. We accessed hitherto unknown physical relations," he said.

He and his research group long studied semiconductor nanomaterials with sizes between 1 nanometer and 10 nm. A major challenge arises when promoting the growth of these materials, as each individual unit grows differently; hence, the spectrum of light emitted by the material

after excitation is broadened, with the various components emitting at slightly different frequencies, and the color of the emission is less precise. When a single particle is isolated, the spectrum becomes narrower, but signal detection is retarded. In other words, spectral resolution is enhanced but at the loss of temporal resolution.

"About five years ago we began working with a technique that can pick out subsets comprising a few thousand identical particles from a set of 1020 nm particles," Padilha said. "This has enabled us to achieve very fine and precise spectral resolution, as well as fine temporal resolution. In this study, we obtained single-particle spectral resolution for a group of particles in an exceptionally short time."

As noted, this experimental solution enabled the researchers to access hitherto unknown physical processes, such as the ultrafast exciton-phonon interaction. It is worth recalling that in condensed matter physics, the phonon is a quasi-particle associated with the quantum of vibration that propagates in a [crystal lattice](#).

There are no immediate technological applications for the results obtained, but in the not-too-distant future, knowledge of physical interactions on the femtosecond timescale can help scientists control the structure of materials such that excitons retain energy from electrical or light impulses for longer periods, retarding decoherence in quantum systems.

"Prolonging coherence is key to the success of devices such as optical switches and single-photon emitters," Almeida said. "Actually, what you aim to do is reduce energy waste to a minimum. When the material changes color, it means it's losing energy. We discovered that this loss is extremely fast. That's what we want to delay."

More information: A. Liu et al, Non-Markovian Exciton-Phonon

Interactions in Core-Shell Colloidal Quantum Dots at Femtosecond Timescales, *Physical Review Letters* (2019). [DOI: 10.1103/physrevlett.123.057403](https://doi.org/10.1103/physrevlett.123.057403)

Provided by FAPESP

Citation: Ultrafast particle interactions could make quantum information devices feasible (2019, October 17) retrieved 22 September 2024 from <https://phys.org/news/2019-10-ultrafast-particle-interactions-quantum-devices.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.