


# Infrasound from underwater volcano erupting shows formation of gigantic bubbles

21 October 2019, by Bob Yirka



WorldView satellite images of the subaerial portion of the vent region of Bogoslof near times when the bubble signals were recorded. Dashed lines circle the approximate submarine vent areas. WorldView data provided under the DigitalGlobe NextView License. Credit: *Nature Geoscience* (2019). DOI: 10.1038/s41561-019-0461-0

A team of researchers with the U.S. Geological Survey and the University of Alaska has found that they could estimate the size of bubbles that form from underwater volcanoes by listening to infrasound produced by bubble formation. In their paper published in the journal *Nature Geoscience*, the group describes their study of the infrasound that was produced by an undersea volcano eruption near an Aleutian island and what they learned from it.

The researchers note that studying undersea volcanoes is challenging due to their remoteness and the unpredictable nature of their eruptions. In this new effort, they report that perseverance and luck allowed them to learn something new about such eruptions—that the bubbles they create are far larger than anyone imagined.

Over the years, there have been accounts by sailors of the sea swelling just prior to the release of gas and particulate from volcanic eruptions below the surface. One such report came from a crewman aboard the *Albatross* during an expedition in the waters around the Aleutian Islands back in 1908. He reported that the ocean swelled into a dome that neared in size to the U.S. Capitol building. Then two years ago, an underwater volcano named Bogoslof erupted. It, too, is part of the Aleutian chain, and is in the process of building the island of Bogoslof. Fortunately, scientists had placed underwater microphones close enough to the [volcano](#) to pick up mysterious infrasounds produced during the [eruption](#).

After studying the low-frequency infrasounds, the researchers found that they were produced by underwater [bubble formation](#). They also found that the bubbles oscillated, indicating that they were changing in size. Further study of the infrasounds allowed the researchers to follow the progression of the bubbles as they made their way to the surface and to measure how big they were. When a single bubble reached the surface, it pushed the water above it into a dome. Then, as the bubble was exposed to the change in pressure, it expanded and contracted several times before finally collapsing. Once it collapsed, the gases and other material in the bubble escaped into the air and created a giant plume. The researchers further report that they measured bubbles as large as 440 meters across, far bigger than the U.S. Capitol dome.

**More information:** John J. Lyons et al. Infrasound from giant bubbles during explosive submarine eruptions, *Nature Geoscience* (2019). DOI: [10.1038/s41561-019-0461-0](https://doi.org/10.1038/s41561-019-0461-0)

© 2019 Science X Network

APA citation: Infrasound from underwater volcano erupting shows formation of gigantic bubbles (2019, October 21) retrieved 24 November 2020 from <https://phys.org/news/2019-10-infrasound-underwater-volcano-erupting-formation.html>

*This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.*