


The forests of the Amazon are an important carbon sink

11 November 2019



The world's tropical forests store huge quantities of carbon in their biomass and thus constitute an important carbon sink. Credit: R.M. Nunes/Shutterstock


The world's tropical forests store huge quantities of carbon in their biomass and thus constitute an important carbon sink. However, current estimates of the amount of carbon dioxide stored in tropical forests of the Amazon vary largely. Scientists at the Helmholtz Centre for Environmental Research (UFZ) have developed an approach that uses recent satellite data to provide much more precise estimates of the amount of biomass in tropical forests than in the past. This makes it possible to obtain a more exact picture of the consequences of droughts and forest fires for the Amazon, according to their article in *Nature Communications*.

Estimating forest biomass from [satellite measurements](#) is still a challenge, as there exists currently no direct measurement method. State-of-the-art satellites equipped with laser or radar instruments now open up a new range of options: they not only measure the height of the forests around the globe but also the entire structure of these forests. The forest modelling team led by

Prof. Dr. Andreas Huth of the UFZ now combined the measurements from a laser satellite with the FORMIND forest model developed at the UFZ; this model uses climate and soil data to simulate the dynamics of forests and the growth of individual trees with a resolution of up to 20 meters. Over 700,000 such laser datasets were evaluated in this way for the Amazon rainforest. The result: important forest attributes, which are crucial in providing a picture of a forest area, can be estimated much more accurately in future. These forest attributes include the above-ground biomass and the rate of forest growth (i.e. gross primary production). "All in all, the uncertainty surrounding the estimates of the forest parameters decreases by between 20 and 43 percent. The estimate of above-ground biomass, for example, has become 25 percent more accurate," says Andreas Huth, one of the authors of the study. This makes it possible to obtain a significantly more precise assessment of the quantity of carbon stored in the forest. "With our approach, we are able to find out more about the carbon cycle—how much is stored in the tropical rainforest, released or resorbed again each year," adds Dr. Rico Fischer, also author of this study and involved in forest modelling at the UFZ.

The UFZ team will also use the new approach to refine their own studies, which only incorporated the forest height to date but not all the information on forest structure, in their biomass estimates. In 2018, they succeeded in simulating the biomass of all 410 billion trees in the Amazon region for the year 2005 by combining laser data from the ICESat satellite with FORMIND. According to one result provided by this method, a total of 76 billion tonnes of carbon is stored in the Amazon rainforest. "This also enabled us to identify which areas of the Amazon region are carbon sinks or sources of carbon," says Andreas Huth. Overall, the rainforest still constitutes a carbon sink by absorbing around 600 million tonnes per year. There are, however, also local carbon sources, such as when trees die

due to drought or are destroyed by fire.



Forest biomass in the Amazon in 2005. Combining the FORMIND forest model with data from the ICESat satellite made it possible to create a detailed biomass map. According to this map, 76 billion tonnes of carbon is stored in the Amazon forest. The red colouring shows areas with a particularly large amount of biomass. Credit: Rödiger et al., *Global Ecol Biogeogr.* 2017

Combining high-resolution satellite data with the FORMIND model now opens up a range of new options for the forest modeling team at the UFZ. For example, the GEDI (Global Ecosystem Dynamics Investigation) mission launched by NASA has been using a new type of laser instrument at the International Space Station (ISS) to measure the global forest since 2018. This data will be available at the end of this year. This would enable UFZ researchers to make statements at six-monthly intervals on how [land use](#) or global warming—to name just two factors—has changed the quantity of biomass stored in the [tropical forests](#) and where the carbon sinks and sources are located. Up-to-date assessments of the consequences of forest fires, such as those in the Amazon region, are also feasible. "As soon as the NASA measurements are available, we will be able to analyze how much [carbon](#) dioxide was emitted by fires in the Amazon," says Rico Fischer.

Another vision held by UFZ researchers is to

integrate the data provided by other satellites and to blend it with FORMIND. This would further reduce the uncertainties contained in the estimates.

Researchers could also benefit from a new radar [satellite](#) mission proposed and planned by German scientists, the Tandem-L mission. One of the mission's objectives is to measure the structure of forests around the world every week by deploying two radar satellites. This would make it possible to quickly identify changes to the [forest](#) over the short term caused by deforestation, [forest fires](#) or droughts, for example, and thus to quantify much more precisely the consequences of land use and climate change. This, according to Rico Fischer and Andreas Huth, would be a further big step for ecological research.

More information: Edna Rödiger et al. From small-scale forest structure to Amazon-wide carbon estimates, *Nature Communications* (2019). DOI: [10.1038/s41467-019-13063-y](https://doi.org/10.1038/s41467-019-13063-y)

Edna Rödiger et al. The importance of forest structure for carbon fluxes of the Amazon rainforest, *Environmental Research Letters* (2018). DOI: [10.1088/1748-9326/aabc61](https://doi.org/10.1088/1748-9326/aabc61)

Provided by Helmholtz Association of German Research Centres

APA citation: The forests of the Amazon are an important carbon sink (2019, November 11) retrieved 3 March 2021 from <https://phys.org/news/2019-11-forests-amazon-important-carbon.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.