

# Albedo matters for the climate, and forestry can have an impact on it

December 4 2019

---



Credit: Aalto University

A research project conducted by Aalto University found that favoring broadleaved species in boreal forestry is a climate-friendly option when considering the forest albedo. Therefore, forest management actions can directly affect forest albedo, and hence the climate, without decreasing forest productivity.

Vegetation cover and land-use changes alter the surface [albedo](#), or the extent to which incoming solar radiation is reflected back to the atmosphere and outer space. Even though land surface albedo is a

critical variable affecting our climate, it is still among the main uncertainties of the radiation budget in current climate modelling. Aalto University's soon ending project funded by the Academy of Finland examined how changes in boreal forests affect albedo and its seasonal variation.

The connection between the on-going environmental changes in the boreal zone and forest albedo has been unknown. Especially important in terms of climate is to assess how forest management actions and natural changes in forests affect albedo. On the other hand, it is critical also to examine how forest albedo and [forest productivity](#) are interconnected.

## **Seasonal dynamics of forests were measured from space**

The project developed an interdisciplinary methodology for assessing boreal albedo and its seasonality based on forest inventory data, forest reflectance modelling and Earth observation data. Field reference data was collected from Finland, Estonia, Sweden, Russia and Alaska (United States).

In the northern forests, seasonality of albedo is large: the albedo can vary between 7 and 80 percent depending on the season. Especially the [snow cover](#) and the duration of the snow-covered season have an important role in albedo regulation. Besides this, species composition, forest density and forest floor vegetation affect forest albedo.

## **Forestry matters for albedo**

The research results demonstrated that forest management can affect boreal forest albedo, but the local circumstances have to be taken into account. Forest density and species composition influence albedo

differently during snow-covered and snow-free seasons. Therefore, the duration of the snow-covered season has a great impact on how forest management affects the mean annual albedo. For example, in the southern boreal region, increasing the fraction of broadleaved species in coniferous forests and favoring mixed forests can increase forest albedo without compromising productivity. In the northern boreal region, on the other hand, the importance of the tree species is not as obvious, and the albedo can be best increased by decreasing the density of the forests. However, as forest density is closely connected to productivity, it is important to take into account also other aspects of forestry such as carbon sequestration and the economic value of forests.

With the knowledge of consequences of forest management practices on the albedo, best-practice recommendations for [forest](#) management in future climate mitigation policies will become more justified. In terms of climate, favoring mixed forests in the southern boreal zone can be recommended. Also, the importance of albedo for the climate increases towards the north. This is why in the boreal zone, acknowledging also the role of albedo in forestry is an essential [climate](#) action.

Provided by Aalto University

Citation: Albedo matters for the climate, and forestry can have an impact on it (2019, December 4) retrieved 3 May 2024 from <https://phys.org/news/2019-12-albedo-climate-forestry-impact.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.