


Capturing alien comets: Simulating rogue bodies on their journey through the solar system

December 20 2019



Comet Borisov in front of a distant background spiral galaxy imaged by the Hubble Space Telescope on November 16, 2019. Credit: NASA, ESA, and D. Jewitt (UCLA)

There should be interstellar comets hiding in our solar system after making a journey of many light-years. Maybe we have already seen one but believed it was a "normal" comet formed in the solar system, according to Tom Hands, astrophysicist at the University of Zürich and member of the NCCR PlanetS.

Comets have fascinated humanity for centuries. Where do these exotic objects come from? According to the most popular theory proposed by Dutch astronomer Jan Oort, during a very early phase of the [solar system](#)'s formation, the giant planets scattered objects into the outer regions far away from the sun. There, the icy rocks and dust particles formed a kind of cloud. Passing stars could scatter these objects back into the inner solar system, where we observe them as comets. Coming from the Oort Cloud, these long-period comets often need more than 200 years for one orbit around the sun.

"We present a second potential origin for such comets," says Tom Hands, postdoc at the Institute for Computational Science of the University of Zürich: "They [could have been] captured out of interstellar space in the relatively recent past."

Two interstellar visitors made headlines in recent years. In 2017, the first such object was detected, an asteroid-like body named 'Oumuamua. In August 2019 amateur astronomer Gennady Borisov found a [comet](#) that came from [interstellar space](#) and will leave the solar system again. 'Oumuamua and Comet Borisov are both leftovers of planet formation in other solar systems, in the same way our comets and asteroids are

thought to be the leftovers of planet formation in our solar system.

Simulating 400 million objects

In the wake of the discovery of the first two interstellar objects, Tom Hands and Walter Dehnen of the University of Munich, Germany, used [computer simulations](#) to study how interstellar objects could be captured by our solar system. "These stowaways form around distant stars before being flung toward us, making a journey of many light-years before encountering Jupiter and being captured into the solar system," explains Hands. "We simulated 400 million such bodies as they approached the sun and Jupiter." The researchers used realistic velocities for these objects based on data from the GAIA mission, and studied how they interact with Jupiter on their journey through the solar system.

This work was done on the VESTA cluster of the University of Zürich. "We used an advanced code which runs on graphics processing units rather than traditional computer processors to enable us to simulate such a large number of objects in a short time," explains Hands. "The simulations took two days in total using around 70 graphics cards. [It would have taken] roughly 140 days if we had only used one card and much, much longer if we had used a normal desktop computer processor."

The results of the simulations now published in the *Monthly Notices of the Royal Astronomical Society (MNRAS)* reveal that in a small minority of cases, the trajectories of the objects are altered enough by Jupiter that they become bound to the solar system. "Although the capture probability is small, there could be anywhere from a few hundred to hundreds of thousands of these alien comets orbiting the sun," says the astrophysicist.

Captured objects are typically on orbits very similar to those of long-

period comets that humanity has observed for centuries, suggesting that they are hiding in plain sight. "If we could identify one, we would have a real possibility to study the composition of material formed in other solar systems in close detail," says Hands.

ESA has recently selected a mission called Comet Interceptor designed to make a fly-by of a long-period or interstellar comet. The University of Bern will provide the camera system and the mass spectrometer for this mission, and will search for differences between these objects and those comets known to have their origins in our solar system.

In an earlier paper published in May 2019, Hands and colleagues studied how close interactions between stars in their birth cluster affect the comets and asteroids formed around each star. They found that objects may become liberated and left "free-floating" in the galaxy, or alternatively "stolen" by other stars. This led them to suggest that the Oort Cloud might be populated partially by objects that were formed around other [stars](#) but then captured by the sun in its birth cluster billions of years ago. This latest study investigates the capture of free-floating asteroids and comets, which may have been liberated from their parent star by the mechanism demonstrated in the earlier study.

More information: T O Hands et al. Capture of interstellar objects: a source of long-period comets, *Monthly Notices of the Royal Astronomical Society: Letters* (2019). [DOI: 10.1093/mnrasl/slz186](https://doi.org/10.1093/mnrasl/slz186)

Provided by NCCR PlanetS

Citation: Capturing alien comets: Simulating rogue bodies on their journey through the solar system (2019, December 20) retrieved 22 September 2024 from <https://phys.org/news/2019-12-capturing-alien-comets-simulating-rogue.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.