


How many stars eventually collide as black holes? The universe has a budget for that

February 3 2020, by Spencer Turney



Artist's conception shows the collisions of two black holes similar to those detected by LIGO and Virgo gravitational-wave detectors. Image credit: LIGO/Caltech/MIT/Sonoma State. Credit: Aurore Simonnet

Since the breakthrough in gravitational wave astronomy back in 2015, scientists have been able to detect more than a dozen pairs of closely located black holes—known as binary black holes—by their collisions into each other due to gravity. However, scientists still debate how many of these black holes are born from stars, and how they are able to get close enough for a collision within the lifetime of our universe.

Now, a promising new study developed by one Vanderbilt astrophysicist may give us a method for finding the number of available [stars](#) in the history of the universe that collide as [binary black holes](#).

The research, which appears today in *The Astrophysical Journal Letters*, will help future scientists interpret the underlying population of stars and test the formation theories of all colliding black holes across cosmic history.

"Researchers up until now have theorized the formation and existence for pairs of black holes in the universe, but the origins of their predecessors, stars, still remains a mystery," said Vanderbilt astrophysicist and lead author of the study Karan Jani. "With this study, we did a forensic study of colliding black holes using the astrophysical observations that are currently available. In the process, we developed a fundamental constraint, or budget, which tells us about the fraction of stars since the beginning of the universe that are destined to collide as black holes."

Leveraging Einstein's general theory of relativity, which tells us how black holes interact and eventually collide, Jani and co-author Abraham Loeb at Harvard University used LIGO events on record to take an inventory of the universe's time and space resources at any given point. They then developed the constraints accounting for each step in the binary black hole process: the number of available stars in the universe, the process of each star transitioning to an individual black hole, and the

detection of the eventual collision of those black holes—picked up hundreds of millions of years later by LIGO as [gravitational waves](#) emitted by the impact.

"From the current observations, we find that 14 percent of all the massive stars in the universe are destined to collide as black holes. That's remarkable efficiency on nature's part," added Jani. "These added constraints in our framework should help researchers trace the histories of [black holes](#), answering old questions and undoubtedly opening up more exotic scenarios."

More information: Karan Jani et al. Global Stellar Budget for LIGO Black Holes, *The Astrophysical Journal* (2020). [DOI: 10.3847/2041-8213/ab6854](#)

Provided by Vanderbilt University

Citation: How many stars eventually collide as black holes? The universe has a budget for that (2020, February 3) retrieved 24 April 2024 from <https://phys.org/news/2020-02-stars-eventually-collide-black-holes.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.