


Bovine embryo completely regenerates placenta-forming cells

7 February 2020



An early bovine embryo regenerating its TE cells which will later form a large part of the placenta. (Left: intact, Middle: after removal of TE, Right: regenerated). Credit: Kohri N. et al., *Journal of Biological Chemistry*. November 8, 2019

A calf was born from an embryo lacking cells which form a large part of the placenta, providing new insight into the regenerative capacity of mammalian embryos.

Mammalian development starts from a [single cell](#)—a [fertilized egg](#). The egg goes through multiple cell divisions to increase its cell numbers and then starts forming a sphere-like structure with a cavity inside, called the blastocyst. The blastocyst consists of two types of [cells](#), the inner cell mass (ICM) and the trophectoderm (TE), which develop into an embryo proper and a large part of the [placenta](#), respectively.

Scientists led by Manabu Kawahara at Hokkaido University have shown that, since bovine ICM cells can regenerate TE, they are capable of forming both the embryo and placenta. The study was published in the *Journal of Biological Chemistry* and became one of the top 50 most viewed papers from November through December 2019 on the journal's website.

To examine the ICM's capacity to regenerate TE, the researchers cultivated [mouse](#) and bovine blastocysts and removed entire TE from both

blastocysts. They found that both blastocysts regained their sphere-like shapes in 24 hours. However, the regeneration rate to reform the blastocyst was remarkably higher in bovine cells (97%) than mouse cells (57%). The more complete recovery of bovine blastocysts in cell numbers compared to mouse blastocysts suggests the bovine cells have a higher regenerative capacity.



Nanami Kohri of the research team at Hokkaido University. Credit: Hokkaido University

Further experiments revealed abnormal protein expression in the TE of mouse regenerated blastocysts, whereas bovine regenerated blastocysts showed normal gene expressions overall.

To test its developmental abilities, the researchers then transferred the regenerated blastocysts to recipient females. After the embryo-transfer, to their surprise, one of the four cows became pregnant and a female calf was naturally born with an apparently normal placenta. In contrast, none of the more than 100 mouse [embryos](#) transferred to recipients developed to term.

"We will continue to monitor the health of the calf born from the regenerated blastocyst," says Manabu Kawahara. "Our study suggests that we can remove and use a large part of TE for genetic testing to breed cattle with improved qualities. Also, further studies could reveal the mechanism of cell fate decision in mammals and its differences between species."

More information: Nanami Kohri et al.

Trophectoderm regeneration to support full-term development in the inner cell mass isolated from bovine blastocyst, *Journal of Biological Chemistry* (2019). [DOI: 10.1074/jbc.RA119.010746](https://doi.org/10.1074/jbc.RA119.010746)

Provided by Hokkaido University

APA citation: Bovine embryo completely regenerates placenta-forming cells (2020, February 7) retrieved 25 November 2020 from <https://phys.org/news/2020-02-bovine-embryo-regenerates-placenta-forming-cells.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.